

(12)

**EUROPEAN PATENT APPLICATION**

(21) Application number: 84301468.9

(22) Date of filing: 06.03.84

(51) Int. Cl.<sup>3</sup>: **C 12 N 15/00**  
**C 12 P 19/34, C 07 C 103/52**  
**C 12 P 21/00, C 12 N 9/48**  
**C 12 N 9/38**  
**//C12R1/19, C12R1/38, C12R1/07,**  
**C12R1/865**

(30) Priority: 28.03.83 GB 8308483

(43) Date of publication of application:  
 10.10.84 Bulletin 84/41

(84) Designated Contracting States:  
 BE CH DE FR GB IT LI NL SE

(71) Applicant: **The Public Health Laboratory Service Board**  
**61 Colindale Avenue**  
**London NW9 5EQ(GB)**

(72) Inventor: **Atkinson, Anthony**  
**Twigley Mill Corner Winterbourne Gunner**  
**Salisbury Wiltshire SP4 6JJ(GB)**

(72) Inventor: **Minton, Nigel Peter**  
**6 Highbury Avenue**  
**Salisbury Wiltshire SP2 7EX(GB)**

(72) Inventor: **Sherwood, Roger Franklin**  
**35 Attwood Road**  
**Salisbury Wiltshire SP1 3PR(GB)**

(74) Representative: **James, Stephen Richard et al,**  
**Procurement Executive Ministry of Defence Patents**  
**1A(4), Room 2014 Empress State Building**  
**Lillie Road London SW6 1TR(GB)**

(54) A leader sequence to promote the secretion of gene products.

(57) A recombinant DNA transfer vector contains a leader  
 sequence polynucleotide which codes for a signal polypep-  
 tide of formula I,

Met - Arg - Pro - Ser - Ile - His - Arg - Thr -  
 Ala - Ile - Ala - Ala - Val - Leu - Ala - Thr - I  
 Ala - Phe - Val - Ala - Gly - Thr

Preferably the transfer vector is a plasmid. In one preferred embodiment the leader sequence polynucleotide is downstream of and in reading phase with a bacterial or yeast promoter and a ribosome binding site, and upstream of and in reading phase with a structural gene. The structural gene may be, for example, the carboxypeptidase G<sub>2</sub>(CPG<sub>2</sub>) gene from the chromosomal DNA of *Pseudomonas* species strain RS-16. Examples of plasmids containing the leader sequence polynucleotide and the CPG<sub>2</sub> gene are pNM1, pNM111, pNM14, pNM21, pNM22, pNM31, pNM32 and pLEC3.

LEADER SEQUENCE TO PROMOTE THE SECRETION OF GENE PRODUCTS

The present invention relates to fragments of specific deoxyribonucleotide sequences that promote the secretion of gene products from cells and in particular to recombinant DNA transfer vectors that contain these fragments.

5       Recent developments in biochemistry have led to the construction of recombinant DNA transfer vectors in which, transfer vectors, for example plasmids, are made to contain exogeneous DNA. In some cases the recombinant incorporates heterologous DNA that codes for polypeptides that are ordinarily not produced by the organism  
10       susceptible to transformation by the recombinant vehicle.

In its basic outline a method of endowing a micro organism with the ability to synthesise a new protein involves three  
general steps:

- 15       (a) isolation and purification of the specific gene or nucleotide sequences containing the genetically coded information for the amino acid sequence of the desired protein or polypeptide,
- 20       (b) recombination of the isolated gene or nucleotide sequence with an appropriate transfer vector, typically DNA of a bacteriophage or plasmid to form a recombinant transfer vector that codes, in part, for the production of the desired protein or polypeptide,

- (c) transfer of the vector to the appropriate micro organism and selection of a strain of the recipient micro organism containing the desired genetic information.

Provided the gene or nucleotide sequence expresses its  
5 protein or polypeptide in the chosen micro organism, growth of the micro organism should then produce the desired protein or polypeptide in significant quantities.

Once the micro organism has been cultured, the protein or polypeptide must be isolated from the undesired materials. This  
10 step is considerably facilitated if the majority of the desired protein or polypeptide is present in the culture medium and/or the periplasmic space of the micro organism. In other words purification may be performed in a more efficient manner if, once expressed, the protein or polypeptide passes through the cell  
15 membrane and out of the cytoplasm.

The passage of the protein or polypeptide through the cell membrane is desirable for two main reasons. First the desired protein or polypeptide will generally be foreign to the micro organism in which it is expressed. In many cases, therefore, it  
20 will be quickly broken down by proteolytic enzymes etc in the cells cytoplasm and will, subsequently, have a short half life within the cell. By transferring the protein or polypeptide out of the cytoplasm soon after expression the stability of the protein or polypeptide will be greatly increased. Second the number of unwanted  
25 genetic materials and products (from which the desired protein or polypeptide must be isolated) will be far greater in the cell's cytoplasm than in the culture medium and/or in the cell's periplasmic space. It can be seen that on both of the above counts the transfer of the protein or polypeptide through the cell membrane  
30 and out of the cytoplasm will greatly facilitate protein or polypeptide isolation.

One way in which the secretion of gene products from the cell's cytoplasm may be promoted is to produce, within the cytoplasm, a preprotein or prepolyptide in which the desired protein  
35 or polypeptide is preceded by a signal polypeptide. The predominantly hydrophobic signal polypeptide directs the desired protein or polypeptide to the cell's periplasmic space, where the signal peptide is removed as the desired protein or polypeptide

traverses the cell membrane.

Many of the known signal peptides contain cysteine residues. These residues have been found to react in the cell membrane and thereby inhibit the efficient transfer of the desired gene product  
5 out of the cell.

It is the primary object of the present invention to provide recombinant DNA transfer vectors containing a leader sequence polynucleotide that codes for a signal peptide that is cysteine free. Other objects and advantages of the present invention will become  
10 apparent from the following description thereof.

According to the present invention there is provided a recombinant DNA transfer vector comprising a leader sequence polynucleotide coding for signal polypeptide of formula I,

Met-Arg-Pro-Ser-Ile-His-Arg-Thr-Ala-Ile-Ala-I  
15 Ala-Val-Leu-Ala-Thr-Ala-Phe-Val-Ala-Gly-Thr

The transfer vector may be a bacteriophage or, which is preferred, a plasmid.

Preferably the majority of the codons in the nucleotide sequence are those preferred for the expression of microbial  
20 genomes. Suitable codons are listed in UK 1,568,047 and UK 2007675A, and these publications are therefore incorporated herein by reference.

In one preferred embodiment of the present transfer vector the nucleotide sequence has formula II

25 5'- ATG CCG CCA TCC ATC CAC CGC ACA  
GCC ATC GCC GCC GTG CTG GCC ACC II  
GCC TTC GTG GCG GGC ACC - 3'

The nucleotide sequence coding for the signal polypeptide (the leader sequence polynucleotide) will preferably be downstream of and  
30 in reading phase with a bacterial or a yeast promoter and a prokaryotic ribosome binding site in the transfer vector. Moreover the leader sequence polynucleotide will either be upstream of an insertion site for a structural gene or, which is preferred, will be upstream of and in reading phase with a structural gene coding for a desired protein or polypeptide. Preferably the gene codes for a eukaryotic, particularly a mammalian, protein or polypeptide.

The structural gene may code, for example, for such eukaryotic proteins as human growth hormone, human insulin or human chorionic somatomammotropin. Alternatively it may code for such prokaryotic

proteins as E.coli  $\beta$ -galactosidase or Pseudomonas carboxy peptidase  $G_2$  (CPG<sub>2</sub>) (Carboxypeptidase  $G_2$  is an enzyme, produced by Pseudomonas species strain RS-16, that has application in cancer chemotherapy. It is a  $Zn^{2+}$  containing dimer of 2 x 42,000 daltons and has high  
5 affinities (Km values of  $10^{-5}$  or  $10^{-6}M$ ) for both 5-methyltetrahydrofolate, the predominant circulatory form of folate in mammals and for the folic acid antagonist methotrexate (MTX), which is widely used in cancer chemotherapy. The enzyme may be used directly for the plasma depletion of reduced folates, essential as co-factors in purine  
10 and particularly in pyrimidine biosynthesis. CPG<sub>2</sub> has been shown to inhibit the development of the Walker 256 carcinoma in vivo and to remove MTX from circulation in patients where prolonged exposure to high doses of MTX leads to toxicity).

Examples of transfer vectors according to the present invention  
15 that code for CPG<sub>2</sub> are pNM1, pNM111, pNM14, pNM21, pNM22, pNM31, pNM32 and pLEC3.

The promoter is preferably a high expression bacterial or yeast promoter for the structural gene in a variety of hosts. The particular choice of promoter will depend on the microorganism to be transformed.  
20 For example the transformation of E.coli will generally be effected by a transfer vector in which an E.coli promoter controls the expression of the structural gene. Examples of E.coli promoters are those present in the plasmids pBR 322 and pAT 153. By contrast, the transformation of Pseudomonas species will generally be effected by a transfer  
25 vector in which a Pseudomonas promoter controls the expression of the structural gene. Examples of Pseudomonas promoters are those present in the plasmid pKT 230 or Pseudomonas chromosomal DNA.

In order to express the structural gene the present transfer vector will be transformed into a suitable microorganism. According  
30 to a further aspect of the present invention therefore there is provided a microorganism transformed by a recombinant DNA transfer vector according to this invention. The microorganism will preferably be a bacterium or yeast in which high expression of the structural gene, within the transfer vector, occurs. Depending on the choice of  
35 promoter the microorganism may be a strain chosen from one of the following bacteria E.coli, Pseudomonas and Bacillus or the yeast Saccharomyces cerevisiae.

Having transformed the microorganism, the protein or polypeptide, for which the structural gene codes, may then be expressed by culturing

the transformed microorganism in a culture medium. It is the primary advantage of the present invention that culturing the transformed microorganism affords a preprotein or prepolypeptide in which the desired protein or polypeptide is preceded by the present signal polypeptide. This means that soon after expression the signal polypeptide directs the desired protein or polypeptide to the cell's periplasmic space, where the signal polypeptide is removed as the desired protein or polypeptide traverses the cell membrane. Since the present signal polypeptide is free of cysteine residues the desired gene product will be efficiently secreted through the membrane.

The present transfer vectors may be prepared by any of the methods that are well known in the recombinant DNA art. For example the leader sequence polynucleotide may be synthesised by the modified triester method of K.Itakura et al, JACS, 1975, 97, 7327 or by the improved oligodeoxynucleotide preparation described in UK 2007675A. The disclosure of both of these references is incorporated herein by reference. The synthesised polynucleotide may then be inserted in a transfer vector, preferably a plasmid. In the transfer vector it will preferably be downstream of and in reading phase with a bacterial or a yeast promoter and a prokaryotic ribosome binding site. The leader sequence polynucleotide should also be either upstream of a structural gene insertion site or upstream of and in reading phase with a structural gene.

Alternatively, DNA fragments containing the leader sequence polynucleotide may be obtained from natural sources, in particular from the chromosomal DNA of *Pseudomonas* species strain RS-16. In this particular case a polynucleotide (formula II above) coding for the present signal polypeptide immediately precedes a structural gene coding for CPG<sub>2</sub>. A number of the DNA fragments containing this leader sequence polynucleotide may therefore be recognised by their ability, on insertion into a plasmid and transformation of a microorganism by the resultant recombinant vector, to enable a microorganism to grow on folate. Examples of such recombinant transfer vectors that contain both a polynucleotide coding for the present signal polypeptide (formula II above) and a structural gene coding for CPG<sub>2</sub> are pNM1, pNM111, pNM14, pNM21, pNM22, pNM31, pNM32 and pLEC3. Of course, once a Fol<sup>+</sup> recombinant vector has been obtained in this way it may be subcloned to afford alternative vectors (either Fol<sup>+</sup> or Fol<sup>-</sup>) that also contain a polynucleotide coding for the present signal polypeptide.

Once a suitable DNA fragment has been isolated it may then be inserted in a transfer vector, preferably a plasmid. In the transfer vector the leader sequence polynucleotide on the inserted fragment should be downstream of and in reading phase with a bacterial or a yeast promoter and a prokaryotic ribosome binding site. The leader sequence polynucleotide should also be either upstream of a structural gene insertion site or upstream of and in reading phase with a structural gene.

The structural gene for insertion downstream of and in reading phase with the present leader sequence polynucleotide may be obtained, for example, by the synthetic methods mentioned above (this is particularly useful for the preparation of genes coding for small proteins, such as human growth hormone, insulin and human chorionic somatomammotropin.) Alternatively the structural gene may be prepared from m-RNA by the use of the enzyme reverse transcriptase or may be isolated from natural sources (chromosomal DNA).

An example of the latter method is the isolation of DNA fragments containing a polynucleotide sequence (shown in Table 1) coding for the enzyme CPG<sub>2</sub> (amino acid sequence also shown in Table 1) from Pseudomonas species strain RS-16 chromosomal DNA. Examples of plasmids containing a CPG<sub>2</sub> structural gene, as well as a polynucleotide coding for the present signal polypeptide (formula II above), are pNM1, pNM111, pNM14, pNM21, pNM22, pNM31, pNM32 and pLEC3.

Once prepared or isolated the Leader sequence polynucleotide and the structural gene will be inserted into a transfer vector, preferably a plasmid, to form a recombinant DNA transfer vector according to the present invention. The insertion step or steps will preferably be effected by one of the well known techniques in this art that employ restriction endonucleases, see for example the methods discussed in UK 2090600A, the disclosure of which is incorporated herein by reference. The choice of transfer vector will be determined by the microorganism in which the leader sequence polynucleotide and structural gene are to be expressed. Generally the transfer vector will be a cloning vehicle that is suitable for transforming the chosen micro-organisms and that displays a phenotypical characteristic, such as antibiotic resistance, by which the recombinant transfer vectors may be selected. Thus, if the micro-organism is to be E-coli, then suitable transfer vectors will be the E-coli plasmids pBR322 and pAT153. Alternatively, if the micro-organism is to be Pseudomonas,

TABLE 1

A Polynucleotide Sequence, coding for CPG<sub>2</sub>, isolated from  
*Pseudomonas* species strain RS - 16 chromosomal DNA

	1							
	Met	Arg	Pro	Ser	Ile	His	Arg	Thr
5' -	ATG	CGC	CCA	TCC	ATC	CAC	CGC	ACA
	10							
Ala	Ile	Ala	Ala	Val	Leu	Ala	Thr	Ala
GCC	ATC	GCC	GCC	GTG	CTG	GCC	ACC	GCC
	20							
Phe	Val	Ala	Gly	Thr	Ala	Leu	Ala	Gln
TTC	GTG	GCG	GGC	ACC	GCC	CTG	GCC	CAG
	30							
Lys	Arg	Asp	Asn	Val	Leu	Phe	Gln	Ala
AAG	CGC	GAC	AAC	GTG	CTG	TTC	CAG	GCA
	40							
Ala	Thr	Asp	Glu	Gln	Pro	Ala	Val	Ile
GCT	ACC	GAC	GAG	CAG	CCG	GCC	GTG	ATC
	50							
Lys	Thr	Leu	Glu	Lys	Leu	Val	Asn	Ile
AAG	ACG	CTG	GAG	AAG	CTG	GTC	AAC	ATC
	60							
Glu	Thr	Gly	Thr	Gly	Asp	Ala	Glu	Gly
GAG	ACC	GGC	ACC	GGT	GAC	GCC	GAG	GGC
	70							
Ile	Ala	Ala	Ala	Gly	Asn	Phe	Leu	Glu
ATC	GCC	GCT	GCG	GGC	AAC	TTC	CTC	GAG
	80							
Ala	Glu	Leu	Lys	Asn	Leu	Gly	Phe	Thr
GCC	GAG	CTC	AAG	AAC	CTC	GGC	TTC	ACG
	90							
Val	Thr	Arg	Ser	Lys	Ser	Ala	Gly	Leu
GTC	ACG	CGA	AGC	AAG	TCG	GCC	GGC	CTG
	90							
Val	Val	Gly	Asp	Asn	Ile	Val	Gly	Lys
GTG	GTG	GGC	GAC	AAC	ATC	GTG	GCC	AAG



	100							
Ile	Lys	Gly	Arg	Gly	Gly	Lys	Asn	Leu
ATC	AAG	GGC	CGC	GGC	GGC	AAG	AAC	CTG
		110						
Leu	Leu	Met	Ser	His	Met	Asp	Thr	Val
CTG	CTG	ATG	TCG	CAC	ATG	GAC	ACC	GTC
		120						
Tyr	Leu	Lys	Gly	Ile	Leu	Ala	Lys	Ala
TAC	CTC	AAG	GGC	ATT	CTC	CCG	AAG	GCC
				130				
Pro	Phe	Arg	Val	Glu	Gly	Asp	Lys	Ala
CCG	TTC	CGC	GTC	GAA	GGC	GAC	AAG	GCC
					140			
Tyr	Gly	Pro	Gly	Ile	Ala	Asp	Asp	Lys
TAC	GGC	CCG	GGC	ATC	CCC	GAC	GAC	AAG
						150		
Gly	Gly	Asn	Ala	Val	Ile	Leu	His	Thr
GGC	GGC	AAC	GGG	GTC	ATC	CTG	CAC	ACG
							160	
Leu	Lys	Leu	Leu	Lys	Glu	Tyr	Gly	Val
CTC	AAG	CTG	CTG	AAG	GAA	TAC	GGC	GTG
								170
Arg	Asp	Tyr	Gly	Thr	Ile	Thr	Val	Leu
CGC	GAC	TAC	GGC	ACC	ATC	ACC	GTG	CTG
Phe	Asn	Thr	Asp	Glu	Glu	Lys	Gly	Ser
TTC	AAC	ACC	GAC	GAG	GAA	AAG	GGT	TCC
180								
Phe	Gly	Ser	Arg	Asp	Leu	Ile	Gln	Glu
TTC	GGC	TCG	CGC	GAC	CTG	ATC	CAG	GAA
	190							
Glu	Ala	Lys	Leu	Ala	Asp	Tyr	Val	Leu
GAA	GCC	AAG	CTG	GCC	GAC	TAC	GTG	CTC
		200						
Ser	Phe	Glu	Pro	Thr	Ser	Ala	Gly	Asp
TCC	TTC	GAG	CCC	ACC	AGC	GCA	GGC	GAC

0121352

Glu	Lys	Leu	210 Ser	Leu	Gly	Thr	Ser	Gly
GAA	AAA	CTC	TCG	CTG	GGC	ACC	TCG	GGC
Ile	Ala	Tyr	Val	220 Gln	Val	Asn	Ile	Thr
ATC	GCC	TAC	GTG	CAG	GTC	AAC	ATC	ACC
Gly	Lys	Ala	Ser	His	230 Ala	Gly	Ala	Ala
GGC	AAG	GCC	TCG	CAT	GCC	GGC	GCC	GCG
Pro	Glu	Leu	Gly	Val	Asn	240 Ala	Leu	Val
CCC	GAG	CTG	GGC	GTG	AAC	GCG	CTG	GTC
Glu	Ala	Ser	Asp	Leu	Val	Leu	250 Arg	Thr
GAG	GCT	TCC	GAC	CTC	GTG	CTG	GCG	ACC
Met	Asn	Ile	Asp	Asp	Lys	Ala	Lys	260 Asn
ATG	AAC	ATC	GAC	GAC	AAG	GCG	AAG	AAC
Leu	Arg	Phe	Asn	Trp	Thr	Ile	Ala	Lys
CTG	GCG	TTC	AAC	TGG	ACC	ATC	GCC	AAG
270 Ala	Gly	Asn	Val	Ser	Asn	Ile	Ile	Pro
GCC	GCC	AAC	GTG	TCG	AAC	ATC	ATC	CCC
Ala	280 Ser	Ala	Thr	Leu	Asn	Ala	Asp	Val
GCC	AGC	GCC	ACG	CTG	AAC	GCC	GAC	GTG
Arg	Tyr	290 Ala	Arg	Asn	Glu	Asp	Phe	Asp
GCG	TAC	GCG	GCG	AAC	GAG	GAC	TTC	GAC
Ala	Ala	Met	300 Lys	Thr	Leu	Glu	Glu	Arg
GCC	GCC	ATG	AAG	ACG	CTG	GAA	GAG	GCG
Ala	Gln	Gln	Lys	310 Lys	Leu	Pro	Glu	Ala
GCG	CAG	CAG	AAG	AAG	CTG	CCC	GAG	GCC

0121352

Asp	Val	Lys	Val	Ile	320 Val	Thr	Arg	Gly
GAC	GTG	AAG	GTG	ATC	GTC	ACG	CGC	GGC
Arg	Pro	Ala	Phe	Asn	Ala	330 Gly	Glu	Gly
CGC	CCG	GCC	TTC	AAT	GCC	GGC	GAA	GGC
Gly	Lys	Lys	Leu	Val	Asp	Lys	340 Ala	Val
GGC	AAG	AAG	CTG	GTC	GAC	AAG	GGC	GTG
Ala	Tyr	Tyr	Lys	Glu	Ala	Gly	Gly	350 Thr
GCC	TAC	TAC	AAG	GAA	GCC	GGC	GGC	ACG
Leu	Gly	Val	Glu	Glu	Arg	Thr	Gly	Gly
CTG	GGC	GTG	GAA	GAG	CGC	ACG	GGC	GGC
360 Gly	Thr	Asp	Ala	Ala	Tyr	Ala	Ala	Leu
GGC	ACG	GAC	GGC	GCC	TAC	GCC	GGC	CTC
Ser	370 Gly	Lys	Pro	Val	Ile	Glu	Ser	Leu
TCA	GGC	AAG	CCA	GTG	ATC	GAG	AGC	CTG
Gly	Leu	380 Pro	Gly	Phe	Gly	Tyr	His	Ser
GGC	CTG	CCG	GGC	TTC	GGC	TAC	CAC	AGC
Asp	Lys	Ala	390 Glu	Tyr	Val	Asp	Ile	Ser
GAC	AAG	GCC	GAG	TAC	GTG	GAC	ATC	AGC
Ala	Ile	Pro	Arg	400 Arg	Leu	Tyr	Met	Ala
GCG	ATT	CCG	CGC	GCG	CTG	TAC	ATG	GCT
Ala	Arg	Leu	Ile	Met	410 Asp	Leu	Gly	Ala
CGC	CGC	CTG	ATC	ATG	GAT	CTG	GGC	GCC
Gly	Lys							
GGC	AAG	TGA - 3'						

Amino acids 1 to 22 are the present signal polypeptide

Amino acids 23 to 415 are the CPG<sub>2</sub> structural gene

NB The leader sequence polynucleotide is the preferred polynucleotide of formula II.

then a suitable transfer vector will be *Pseudomonas* pKT230.

The present recombinant DNA transfer vectors, micro-organisms transformed by the present recombinant DNA transfer vectors and processes for the preparation of said vectors and micro-organisms will now be described by way of example only, with particular reference to the Figures in which:

- Figure 1 is a restriction enzyme cleavage site map of pNM1,  
 Figure 2 is a restriction enzyme cleavage site map of pNM111,  
 Figure 3 is a restriction enzyme cleavage site map of pNM14,  
 Figure 4 is a restriction enzyme cleavage site map of pNM21,  
 Figure 5 is a restriction enzyme cleavage site map of pNM22, and  
 Figure 6 illustrates the process for the preparation of a recombinant plasmid containing both the present leader sequence polynucleotide and the  $\beta$ -Galactosidase structural gene, and  
 Figure 7 is a restriction enzyme cleavage site map of pLEC3.

#### Materials and Methods

##### Bacterial strains and plasmids

The bacterial strains used were *Escherichia coli* W5445 (pro leu thi thr<sup>o</sup> sup E44 lac Y ton A r<sup>-</sup> m<sup>-</sup> Str<sup>R</sup>) *Pseudomonas putida* 2440 (r<sup>-</sup>) and *Pseudomonas* sp strain RS-16. The plasmids employed were pBR322 (F Bolivar et al Gene, 1977, 2, 95), pAT153 (A J Twigg et al, Nature, 1980, 283, 216) and pKT230 (M Bagdasarín et al, Gene 1981, 16, 237) and pROG5 (R.F.Sherwood et al, The Molecular Biology of Yeast, 1979 Cold Spring Harbor Publications).

##### Media and culture conditions

*E.coli* was routinely cultured in L-broth (1% tryptone, 0.5% yeast extract, 0.5% NaCl). Solidified medium (L-agar) consisted of L-broth with the addition of 2% (w/v) agar (Bacto-Difco). Antibiotic concentrations used for the selection of transformants were 50  $\mu$ g/ml ampicillin, 15  $\mu$ g/ml tetracycline and 30  $\mu$ g/ml kanamycin. In the case of *E.coli* these were conducted in 2YT liquid medium (1.6% tryptone, 1% yeast extract, 0.5% NaCl) containing 1% glucose, and 0.05% folate where appropriate. The pseudomonads were grown in a minimal salts solution consisting of per litre: MgSO<sub>4</sub>, 0.05g; CaCl<sub>2</sub>, 2H<sub>2</sub>O, 0.05g; FeSO<sub>4</sub>.7H<sub>2</sub>O, 0.005g; MnSO<sub>4</sub>, 0.0015g; Na<sub>2</sub>Mbo<sub>4</sub>, 2H<sub>2</sub>O, 0.0015g; KH<sub>2</sub>PO<sub>4</sub>; 5g; K<sub>2</sub>HPO<sub>4</sub>.3H<sub>2</sub>O, 12g; glutamate, 10g. The minimal medium employed for *E.coli* was M9 medium (J Miller, Experiments in molecular genetics, Cold Spring Harbor Laboratory, Cold Spring Harbor, New York, 1972).

##### Purification of DNA

Plasmids were purified from chloramphenicol amplified cultures (D B Clewell, J Bacteriol, 1972, 110, 667) by Brij-lysis (D B Clewell et al, Proc Natl Acad Sci, USA, 1969, 62, 1159) and subsequent caesium

chloride-ethidium bromide density gradient centrifugation (A Colman et al, Eur.J Biochem, 1978, 91, 303). A rapid, small scale plasmid isolation technique (Burnboim et al, Nuc.Acids Res, 1979, 7, 1513) was also employed for screening purposes. Chromosomal DNA from the donor  
5 Pseudomonad strain (RS-16) was prepared essentially as described by J Marmar, J.Mol.Biol, 1961, 3, 208.

Restriction, ligation and transformation methods

Restriction endonucleases and DNA ligase were purchased from Bethesda Research Laboratories and used in the buffers and under the conditions  
10 recommended by the supplier. Transformation of E.coli was essentially as described by S N Cohen et al., Proc.Natl.Acad.Sci., USA, 1972, 69, 2110, while Ps.putida was transformed by the method of M Bagdasarian and K N Timmis, Current Topics in Microbiology and Immunology, Eds P H Hofschneider and W Goebel, Springer Verlag, Berlin, 1981, p 47.

15 Agarose gel electrophoresis

Digests were electrophoresed in 0.8% agarose slab gels (10 cm x 20 cm x 0.5 cm) on a standard vertical system (Raven), employing Tris-borate-EDTA buffer. Electrophoresis of undigested DNA was at 125V, 50 mA for  
20 3 hours, while digested DNA was electrophoresed at 15V, 10 mA for 16 hours. Fragment sizes were estimated by comparison with fragments of  $\lambda$  DNA digested with HindIII and  $\lambda$  DNA cut with both HindIII and EcoRI. Fragments were isolated from gels using electroelution (M W McDonnell et al, Proc. Natl.Acad.Sci, USA, 1977, 74, 4835).

Determination of carboxypeptidase G<sub>2</sub> activity

25 Bacteria were grown in 1 litre batch culture and 100 ml samples taken at various stages in the growth phase. Samples were cooled on ice, centrifuged at 13,000 x g for 10 minutes and resuspended and frozen in 5 ml of 0.1 M Tris HCl, pH 7.3 containing 0.2 mM ZnSO<sub>4</sub>. The cells were disrupted using a MSE Ultrasonic Disintegrator (150 W) at medium  
30 frequency, amplitude 2, for three 30-second intervals on ice. Cell debris was removed by centrifugation at 10,000 x g for 5 minutes. CPG<sub>2</sub> activity was determined after J L McCullough et al, J.Biol.Chem, 1971, 246, 7207. A 1ml reaction cuvette containing 0.9 ml of 0.1 M Tris-HCl, pH 7.3 plus 0.2 mM ZnSO<sub>4</sub> and 0.1 ml of 0.6 mM methotrexate was  
35 equilibrated at 37°C. Enzyme extract was added to the test cuvette and the decrease in absorbance at 320 nm measured using a Pye-Unicam SP1800 double-beam spectrophotometer. Enzyme activity per ml extract

was calculated as  $\Delta$  320 nm absorbance/min divided by 8.3, which is equivalent to the hydrolysis of 1  $\mu$ mol of MEX/min at 37°C. Protein concentration was determined by the method of M M Bradford, Anal Biochem, 1976, 72, 248.

5 Cell fractionation techniques

- Bacterial cultures were grown in the low phosphate medium of H C Neu and L A Heppel, (J Biol Chem, 1964, 240, 3685), supplemented with 100  $\mu$ g/ml ampicillin, to an OD<sub>450</sub> = 1.0. 40 ml of culture was centrifuged at 5000 g for 10 min, washed in 5 ml of 10 mM Tris-HCl pH 7.0, and resuspended in 0.9 ml 0.58 M sucrose, 0.2 mM DTT, 30 mM Tris-HCl pH 8.0. Conversion to spheroplasts was achieved by the addition of 20  $\mu$ l of lysozyme (2 mg/ml), 40  $\mu$ l 0.1 M EDTA, and incubation at 23°C for 10 min (HC Neu et al, J Biol Chem, 1964, 239, 3893). The spheroplasts were placed on ice and 0.1 ml of 30% (w/v) BSA added, followed by 5 ml of sucrose-tris buffer. Sedimentation of the spheroplasts was achieved by centrifugation at 5000g for 10 min and the supernatant retained as the 'periplasmic' fraction. The pellet was resuspended in 5 ml 10 mM Tris-HCl, 0.2 mM DTT pH 7.0 and and sonicated at 20 Kc/sec, 2 Amps for 15 sec. Remaining whole cells were removed by centrifugation at 1000 x g for 10 min. Centrifugation at 100000 x g for 1 hr, at 4°C, separated the soluble (cytoplasmic) proteins from the particulate (membrane-bound) proteins. The membrane pellet was resuspended in 1 ml of 10mM Tris-HCl, 0.2 mM DTT, pH 7.0.
- 25 CPG<sub>2</sub> was assayed as described. Alkaline phosphatase was assayed according to J Miller, Experiments in Molecular genetics, Cold Spring Harbor Laboratory, Cold Spring Harbor, New York, 1972, NADH oxidase according to M J Osborn et al, j Biol Chem, 1972, 247, 3962 and glyceraldehyde - 3 - phosphate dehydrogenase after K Suzuki et al, FEBS, 1971, 13, 217.
- 30

Example 1

Preparation of recombinant plasmid pNML (A plasmid containing both the present leader sequence polynucleotide and the CPG<sub>2</sub> structural gene)

- 5 To isolate the gene for carboxypeptidase G<sub>2</sub> together with the leader sequence polynucleotide chromosomal DNA prepared from the Pseudomonas host (strain RS-16) was partially digested with Sau3A and fragments of between 6-8 Md isolated from agarose gels by electroelution. The 'sized' DNA was ligated with alkaline phosphatase treated BamHI cut pBR322, transformed into E.coli W5445, and Ap<sup>r</sup> transformants selected. Of the 3,500 Ap<sup>r</sup> colonies obtained, approximately 70% were TC<sup>r</sup>. Utilisation of a rapid plasmid isolation technique on 50 Ap<sup>r</sup> TC<sup>r</sup> transformants demonstrated that 90% of the
- 10 check on the authenticity of the gene bank, the individual clones were screened for the acquisition of a Leu<sup>+</sup> phenotype. Two such clones were identified. Both carried a plasmid capable of transforming leuB (B-isopropylmalate dehydrogenase) E.coli mutants to prototrophy.
- 15 Acquisition of a functional CPG<sub>2</sub> gene should enable E.coli to utilise folic acid as a carbon source. The 2,400 gene bank clones were screened for the ability to grow on minimal medium containing folate as the sole source of carbon (ie Fol<sup>+</sup>). A single Fol<sup>+</sup> clone was detected and shown to harbour a plasmid
- 25 capable of transforming plasmid-minus W5445 to the Fol<sup>+</sup> phenotype. Classical restriction mapping of this plasmid (pNML) was undertaken which revealed the presence of a 5.9 Md insert of pseudomonad DNA within pBR322. The restriction enzyme cleavage site map of pNML is given in Figure 1. The nucleotide sequence of the
- 30 leader sequence polynucleotide and the CPG<sub>2</sub> structural gene is given in Table 1.

Example 2

Subcloning of plasmid pNML to form pNML11

- 35 In order to pinpoint the position of the CPG<sub>2</sub> gene and the leader sequence polynucleotide within the 5.9 Md insert, subcloning of various restriction enzyme fragments, into pBR322, was undertaken. A functional CPG<sub>2</sub> gene was shown not to occur on



XhoI or SphI fragments of the pNM1 insert, but was present on a 3.1 Md BglII fragment. This latter fragment was cloned into the BamHI site of pBR322 to give pNM11 (6.0 Md). A further reduction in the size of pNM11 was achieved by digesting with Sall and

5 religating the resultant fragment to yield pNM111. In addition, plasmids in which the smaller 0.95 Md Sall fragment had become inserted in the opposite orientation to the parent plasmid (pNM11) were  $Fol^-$ . Taken together these subcloning results indicate that

10 the BglII site at 4.14 and the Sall site at 6.03 on pNM1. Furthermore, the gene contains a SphI (5.17), Sall (5.07) and at least one XhoI (4.56 and/or 5.56) site. The restriction enzyme cleavage site map of pNM111 is given in Figure 2.

### Example 3

- 15 Preparation of recombinant plasmid pNM14. (A plasmid containing both the present leader sequence polynucleotide and the  $CPG_2$  structural gene)

The 3.1 Md Bgl II fragment from Example 2 above was partially digested with Sau3A. These fragments were then cloned into the

20 Bam HI site of pAT153 and transformed into E coli W5445. Of the two  $Ap^r$   $Tc^s$   $Fol^+$  colonies obtained, one contained a plasmid which had acquired an extra Sal I and Bam HI site, this was pNM 14. The restriction enzyme cleavage site map of pNM 14 is given in Figure 3. Sequencing of the leader sequence polynucleotide and

25 the  $CPG_2$  structural gene present in pNM 14 gave the nucleotide structure shown in Table 1. DNA sequencing of pNM 14 also revealed that the Sal I - Bam HI fragment was a duplication of a segment of DNA from within the insert (marked \* on Figure 3) composed of two contiguous Sau 3A fragments.

### 30 Example 4 and 5

Preparation of recombinant plasmids pNM 21 and pNM 22 (Plasmids containing both the present leader sequence polynucleotide and the  $CPG_2$  structural gene)

- The 3.1 Md Bgl II fragment from Example 2 was cloned into the
- 35 Bam HI site of pAT 153 and transformed into E coli W 5445. Two  $Ap^r$   $Tc^s$   $Fol^+$  colonies were obtained, one containing a plasmid pNM 21 in which the fragment was inserted in the opposite orientation to

pNM1 and one containing a plasmid pNM22 in which the fragment was inserted in the same orientation as pNM1. The restriction enzyme cleavage site maps of pNM21 and pNM22 are given in Figures 4 and 5 respectively.

5       The two plasmids, pNM21 and pNM22 both transformed E.coli to Fol<sup>+</sup>, indicating that a pseudomonad promoter was present on the 3.1Md fragment. However, cells carrying the plasmid pNM21, in which the BglIII fragment was cloned in the opposite orientation to pNM1, exhibited more rapid growth with folic acid as the sole  
10 carbon source. This difference was clearly visible on agar medium, where colonies developed concentric yellow 'halos' of precipitated pteronic acid, the insoluble product of folate hydrolysis.

Confirmation that pNM21 gave enhanced expression of CPG<sub>2</sub> over pNM22, was obtained by assaying enzyme production during  
15 batch growth of cells containing either plasmid. (The cells were grown in complex medium supplemented with 1% (w/v) glucose and where appropriate 0.05% (w/v) folic acid. The generation time was 56-66 min. The culture was sampled at hourly intervals and whole cells were disrupted by sonication. Enzyme activity was  
20 determined in the centrifugal extract). Results are given in Table 2.

The expression of CPG<sub>2</sub> from the plasmids pNM22 and pNM1 was 2.5 units/litre of culture, representing 0.005% soluble protein. In contrast, expression from pNM21 was 3000-3500 units/litre of  
25 culture, which represented 4.7% soluble protein. As the cloned gene is inserted into the BamHI site of pAT153, the observed higher expression of pNM21 is almost certainly due to transcriptional read through from the Tc promoter. The low expression of CPG<sub>2</sub> carried on plasmids pNM1 and pNM22 is consistent with the  
30 view that Pseudomonas promoters function poorly in E.coli. It is also apparent from Table 2 that in the presence of folate there is a two-fold increase in the specific activity of enzyme measured in cell sonicates. This phenomenon has been observed in all experiments, but does not seem to be associated with  
35 classical induction of the CPG<sub>2</sub> gene, as overall enzyme yield in the presence or absence of folate remains at about 3000 u/litre culture. It in fact reflects a consistent depression in the

soluble protein levels measured in sonicates from cells grown in the presence of folate. There is no obvious difference in growth rate of cells grown with folate and the reasons for this result are not clear.

5 TABLE 2: CARBOXYPEPTIDASE G PRODUCTION BY *E. COLI* W5445  
 , CONTAINING THE PLASMIDS pNM1, pNM21 and pNM22.

CULTURE		CARBOXYPEPTIDASE G <sub>2</sub> SPECIFIC ACTIVITY (U/MG SOLUBLE PROTEIN)					
AGE (HR)		pNM1		pNM22		pNM21	
		-FOL	+FOL	-FOL	+FOL	-FOL	+FOL
10	1	-	-	-	-	11.5	13.4
	2	-	-	-	-	12.9	9.6
	3	.008	.005	.010	.019	13.9	23.3
	4	.009	.011	.015	.013	12.3	26.9
	5	.007	.019	.016	.016	11.5	25.6
15	6	.005	.024	.014	.023	13.7	24.1
	7	.015	.029	.024	.043	13.2	20.6
	8	.013	.028	.024	.046	13.0	23.6

#### Expression of the cloned gene in *Ps. putida*

20 The observation that the CPG<sub>2</sub> gene was expressed in *E. coli* regardless of the orientation of the gene within the vector suggested that the promoter region of the CPG<sub>2</sub> gene had been cloned with the structural gene and the leader sequence polynucleotide. The low expression of CPG<sub>2</sub> within *E. coli* from its natural promoter (pNM1, pNM22, pNM111) confirmed other findings that *Pseudomonas*  
 25 promoters are poorly recognised by *E. coli* RNA polymerases. It would be expected that if the gene was introduced back into a *pseudomonad* cellular environment, then improved expression from the *Pseudomonas* promoter should result. The 3.1 kb *Bgl*III fragment was subcloned into the *Pseudomonas* cloning vector pKT230 at its single *Bam*HI site.

Two plasmids were obtained, pNM31 and pNM32 representing the two possible orientations of the cloned gene. These plasmids were transformed into Ps.putida 2440 by the method of Bagdasarian and Timmis. Pseudomonad cells carrying both plasmids were cultured in minimal salts medium and enzyme production monitored.

Yields of 500-1000 units/litre of culture were obtained regardless of gene orientation within the plasmid. Specific activity of the enzyme in cell sonicates was 1.5 to 4.0 U/mg protein representing 0.3 to 0.7% soluble protein (compared with < 0.05% soluble protein in the donor strain RS-16). This result strongly indicates that the CPG<sub>2</sub> promoter is present and operating in a pseudomonad background. When the same plasmids were transformed into E.coli W5445 12-40 U<sub>n</sub>its/litre were found at specific activity < 0.07 U/mg (< 0.01% soluble protein).

#### 15 Periplasmic localisation of CPG<sub>2</sub>

There is evidence that CPG<sub>2</sub> is located in or near the periplasmic space of Pseudomonas strain RS-16. Pterioic acid, the product of CPG hydrolysis of folic acid is extremely insoluble and is found predominantly outside the cell in both liquid and solid media. Exogenous pterioic acid is also seen in E.coli cultures containing the cloned gene when folic acid is present in the medium. This is clearly demonstrated by the 'halo' of precipitated pterioic acid observed around colonies carrying plasmids in which expression of CPG<sub>2</sub> is from the Tc promoter of pBR322 (eg pNM21).

25 The localisation of CPG<sub>2</sub> produced by E.coli cells carrying pNM21 was examined by the separation of cellular proteins into cytoplasmic, periplasmic, and whole membrane fractions. As a control, levels of three marker enzymes, alkaline phosphatase (periplasmic), glyceraldehyde-3-phosphate dehydrogenase (cytoplasmic) and NADH.O<sub>2</sub> oxidoreductase (membrane-bound), were also determined. As can be seen from Table 3 97% of the CPG<sub>2</sub> activity occurs in the periplasm, equivalent to the marker periplasmic enzyme, alkaline phosphatase. This confirms the presence in pNM21 of a leader sequence polynucleotide next to the CPG<sub>2</sub> gene that codes for a signal polypeptide according to this invention that promotes the secretion of CPG<sub>2</sub> from the cytoplasm into the periplasmic space.

Carboxypeptidase G<sub>2</sub> synthesised in E.coli

The specific activity of CPG<sub>2</sub> in crude cell extracts of cells carrying pNM21 was 50-fold higher than equivalent extracts from Pseudomonas strain RS-16. To determine whether the cloned gene=

5 product in E.coli had the same properties as CPG<sub>2</sub> from the pseudomonad, enzyme was purified from E.coli carrying pNM21. The specific activity of purified CPG<sub>2</sub> (single band SDS-PAGE) was 535 U/mg of protein, which compares to 550 U/mg of protein from the

10 pseudomonad. CPG<sub>2</sub> purified from E.coli clone pNM21 co-chromatographed with CPG<sub>2</sub> from Pseudomonas strain RS-16 at a sub-unit molecular weight value of 42,000 daltons. Km values using methotrexate as substrate were  $7.4 \times 10^{-6} \text{M}$  and  $8.0 \times 10^{-6} \text{M}$  respectively. In addition, antiserum raised against the Pseudomonas enzyme

15 indicated immunological identity between the E.coli and Pseudomonas CPG<sub>2</sub>, as a confluent precipitation line was formed on Ouchterlony double diffusion analysis.

TABLE 3

## Localisation of Carboxypeptidase

FRACTION	CPG <sub>2</sub>	ENZYME ACTIVITY		
		AP	GAPDH	NADHOX
Periplasmic	97.0	97.1	6.8	0.25
Cytoplasmic	2.6	2.3	93	8.4
Membrane-bound	0.4	0.6	0.2	89.1

AP = Alkaline phosphatase

GAPDH = Glyceraldehyde-3-phosphate dehydrogenase

NADHOX = NADH.O<sub>2</sub> oxidoreductase

Example 6Preparation of a recombinant plasmid containing both the present leader sequence polynucleotide and the B-Galactosidase structural gene

5        Plasmid pNM14 (Example 3) was treated with Sau 3A (GATC) and the fragments were cloned into the Bam HI site of M13 mp7 template DNA (single stranded DNA (Step A of Figure 6). The product carrying a 318bp Sau 3A fragment coding for the present signal polypeptide and the first 22 amino acids of CPG<sub>2</sub> (nucleotide sequence of this  
10       fragment shown in Table 4) was selected and made double stranded. The DNA coding for the signal polypeptide (and the first part of CPG<sub>2</sub>) was then excised as an Eco RI fragment. This Eco RI fragment was then cloned into the promoter cloning vector E.coli pMC1403 (M.J. Casadaban et al, J Bacteriol, 1980, 143, 971), which carries  
15       only the structural gene (lac Z) for B-galactosidase (ie no promoter and no ATG start codon) (Steps B and C of Figure 6). Plasmids were obtained in which the Eco RI fragment had inserted in both orientations, however, only those in which fusion of the CPG<sub>2</sub> sequence to the B-galactosidase sequence had occurred (i) yielded  
20       a 0.34 Kb fragment upon digestion with BamHI; (ii) enabled the host cell to hydrolyse the colourless lactose analogue, BCIG, and impart a blue colouration to colonies. The 0.34 Kb BamHI fragment has been recloned into M13mp7 and sequenced to confirm that fusion has  
25       occurred. The 'precursor' fusion produced will consist of the signal peptide, the first 22 amino acids of CPG<sub>2</sub>, 6 amino acids derived from the M13mp7 and pMC1403linker units, and B-galactosidase from its 8th amino acid onward.

Localisation experiments have been performed on cells carrying a plasmid coding for the 'fusion gene' where the cellular proteins  
30       have been fractionated into periplasmic, cytoplasmic and membrane fractions. In these experiments an organism (E.coli MC 1061) which is deleted for the lac Z gene was grown in phosphate medium (H.C. Neu et al, J Biol Chem, 1964, 240, 3685) and periplasmic enzymes were released from the harvested cells by conversion to spheroplasts.  
35       Separation of soluble proteins (cytoplasmic) from particulate proteins (membrane band) was achieved by sonicating the harvested

spheroplasts and subsequent centrifugation at 100,000g for 1hr, to sediment the cell membrane (T.J.Silhary et al, Proc Natl Acad Sci USA, 1976, 73, 3423).

5 The results given in Table 5 demonstrate the presence of 50% of the B-galactosidase activity in the periplasmic space. This result is in direct contrast to similar work involving fusion of other periplasmic protein signal sequences to B-galactosidase, where the fusion proteins are not exported, but become jammed in the membrane (P.J. Bassford et al, J Bacteriol, 1979, 139, 19 and  
10 S D Emr et al, J Cell, Biol, 1980, 86, 701).

TABLE 4

The Polynucleotide Sequence of the 318 bp Sau 3A Fragment from  
Recombinant Plasmid pNM14

5' - G	ATC	CAC	GCA	CTG	AAG	GCG	CGC	GGC
AAG	ACG	CGC	GGC	GTG	GCG	ACG	CTG	TGC
ATC	GGC	GGG	GGC	GAA	GGC	ACC	GCA	GTG
GCA	CTC	GAT	TGC	TAT	AAG	AAC	CAT	GGC
TGG	GGA	CGC	CCG	ACA	ACA	GGC	GTC	CAC
CAG	CTT	TTT	TCA	TTC	CGA	CAA	CCC	GAA
CGA	ACA	ATG	CGT	AGA	GCA	GGA	GAT	TCC

Table 4 (comtd)

	Met	Arg	Pro	Ser	Ile	His	Arg	Thr
	ATG	CGC	CCA	TCC	ATC	CAC	CGC	ACA
	Ala	Ile	Ala	Ala	Val	Leu	Ala	Thr
	GCC	ATC	GCC	GCC	GTG	CTG	GCC	ACC
5	Phe	Val	Ala	Gly	Thr	Ala	Leu	Ala
	TTC	GTG	GCG	GGC	ACC	GCC	CTG	GCC
	Lys	Arg	Asp	Asn	Val	Leu	Phe	Gln
	AAG	CGC	GAC	AAC	GTG	CTG	TTC	CAG
	Ala	Thr	Asp	Glu	Gln	Pro	Ala	Val
10	GCT	ACC	GAC	GAG	CAG	CCG	GCC	GTG
								ATC

NB. This fragment carries the leader sequence coding for the signal polypeptide, a part of the CPG<sub>2</sub> structural gene coding for the first 22 amino acids of the protein, the ATG start codon, the CPG<sub>2</sub> ribosome binding site (AGGA.) and other components of the CPG<sub>2</sub> promoter region.

TABLE 5

Localisation of Signal Peptide - B-galactosidase Fusion Protein

	CPG <sub>2</sub> /B-GAL	% LOCALISATION <sup>a</sup>		
		AP	GAPDH	NADHOX
Periplasmic	50.3	97.3	3.4	0.4
Cytoplasmic	30.9	2.5	95.3	8.2
Membrane-bound	18.8	0.2	1.3	89.4

- <sup>a</sup> = average results from 4 experiments
- CPG<sub>2</sub>/B-GAL = Carboxypeptidase G<sub>2</sub>-B-galactosidase fusion protein
- AP = Alkaline phosphatase
- GAPDH = Glyceraldehyde-3-phosphate dehydrogenase
- NADHOX = NADH.O<sub>2</sub> Oxidoreductase



Example 7

Preparation of a recombinant plasmid, containing both the present leader sequence polynucleotide and the CPG<sub>2</sub> structural gene, able to replicate in E.coli and S.cerevisiae

- 5 A 2.03 kilobase BamHI fragment coding for the present signal polypeptide and the entire CPG<sub>2</sub> molecule was cloned in both orientations into the BamHI site of an E. coli/S. cerevisiae shuttle vector pROG5 (R.F. Sherwood and R.K. Gibson, The Molecular Biology of Yeast, 1979, Cold Spring Harbor Publications) to give plasmids pLEC3 and
- 10 pLEC4 (Figure 7). These plasmids were transformed into S. cerevisiae strain LL20 by the lithium acetate induced transformation method described by Ito et al., J. Bact., 1983, 153, 163. Yields equivalent to 10-20 units/litre of culture volume were obtained regardless of gene orientation within the plasmid. Specific activity of the enzyme
- 15 in total cell extracts was 0.2-0.3u/mg protein representing 0.005% soluble protein. This level of expression from the pseudomonad promotor in a yeast background is similar to the level found when the gene was reading from its own promotor in E.coli ( 0.01% soluble protein).

- Localisation experiments have been performed on yeast cells
- 20 carrying the above plasmids by sphaeroplasting the cells using standard techniques described by J.B.D. Beggs, Nature, 1978, 275, 105. Periplasmic enzymes, localised outside of the cell membrane, were released when the cell wall was removed. The osmotic stabiliser (1.2M sorbitol) was then replaced by 0.1M Tris-HCl buffer, pH 7.3 containing
- 25 0.2mM ZnCl<sub>2</sub> to lyse the sphaeroplasts and the whole centrifuged at 100,000 x g for 1 hour to separate proteins in the soluble cytoplasmic fraction from membrane bound proteins. The results in Table 6 demonstrate the presence of 64% of the CPG<sub>2</sub> activity in the periplasmic fraction and a further 16% associated with the cell membrane
- 30 fraction.

TABLE 6

Localisation of CPG<sub>2</sub> in S. cerevisiae

	<u>% CPG<sub>2</sub> activity</u>
Periplasmic	64
Cytoplasmic	20
Membrane bound	16

CLAIMS

1. A recombinant DNA transfer vector comprising a leader sequence polynucleotide characterised in that the leader sequence polynucleotide codes for a signal polypeptide of formula I,  
 5       Met - Arg - Pro - Ser - Ile - His - Arg - Thr -  
           Ala - Ile - Ala - Ala - Val - Leu - Ala - Thr -       I  
           Ala - Phe - Val - Ala - Gly - Thr
2. A recombinant DNA transfer vector according to claim 1 characterised in that the leader sequence polynucleotide is  
 10   of formula II,  
       5' - ATG CGC CCA TCC ATC CAC CGC ACA  
           GCC ATC GCC GCC GTG CTG GCC ACC       II  
           GCC TTC GTG GCG GGC ACC - 3'
3. A recombinant DNA transfer vector according to either  
 15   claim 1 or claim 2 characterised in that the leader sequence polynucleotide is downstream of and in reading phase with a bacterial or a yeast promoter and a prokaryotic ribosome binding site.
4. A recombinant DNA transfer vector according to any one  
 20   of claims 1 to 3 characterised in that the leader sequence polynucleotide is upstream of and in reading phase with a structural gene.
5. A recombinant DNA transfer vector according to claim 4 characterised in that the structural gene codes for human  
 25   growth hormone, human insulin or human chorionic somatomammotropin.
6. A recombinant DNA transfer vector according to claim 4 characterised in that the structural gene codes for E.coli  
        $\beta$  - galactosidase.
7. A recombinant DNA transfer vector according to claim 4 characterised in that the structural gene codes for  
 30   Pseudomonas carboxypeptidase G<sub>2</sub> (CPG<sub>2</sub>).
8. A recombinant DNA transfer vector according to claim 7 comprising a polynucleotide of formula


	1							
	Met	Arg	Pro	Ser	Ile	His	Arg	Thr
5' -	ATG	CGC	CCA	TCC	ATC	CAC	CGC	ACA
	10							
Ala	Ile	Ala	Ala	Val	Leu	Ala	Thr	Ala
GCC	ATC	GCC	GCC	GTG	CTG	GCC	ACC	GCC
	20							
Phe	Val	Ala	Gly	Thr	Ala	Leu	Ala	Gln
TTC	GTG	GCG	GGC	ACC	GCC	CTG	GCC	CAG
	30							
Lys	Arg	Asp	Asn	Val	Leu	Phe	Gln	Ala
AAG	CGC	GAC	AAC	GTG	CTG	TTC	CAG	GCA
	40							
Ala	Thr	Asp	Glu	Gln	Pro	Ala	Val	Ile
GCT	ACC	GAC	GAG	CAG	CCG	GCC	GTG	ATC
	50							
Lys	Thr	Leu	Glu	Lys	Leu	Val	Asn	Ile
AAG	ACG	CTG	GAG	AAG	CTG	GTC	AAC	ATC
	60							
Glu	Thr	Gly	Thr	Gly	Asp	Ala	Glu	Gly
GAG	ACC	GGC	ACC	GGT	GAC	GCC	GAG	GGC
	70							
Ile	Ala	Ala	Ala	Gly	Asn	Phe	Leu	Glu
ATC	GCC	GCT	GCG	GGC	AAC	TTC	CTC	GAG
	80							
Ala	Glu	Leu	Lys	Asn	Leu	Gly	Phe	Thr
GCC	GAG	CTC	AAG	AAC	CTC	GGC	TTC	ACG
Val	Thr	Arg	Ser	Lys	Ser	Ala	Gly	Leu
GTC	ACG	CGA	AGC	AAG	TCG	GCC	GGC	CTG
	90							
Val	Val	Gly	Asp	Asn	Ile	Val	Gly	Lys
GTC	GTG	GGC	GAC	AAC	ATC	GTG	GGC	AAG

	100							
Ile	Lys	Gly	Arg	Gly	Gly	Lys	Asn	Leu
ATC	AAG	GGC	CGC	GGC	GGC	AAG	AAC	CTG
		110						
Leu	Leu	Met	Ser	His	Met	Asp	Thr	Val
CTG	CTG	ATG	TCG	CAC	ATG	GAC	ACC	GTC
		120						
Tyr	Leu	Lys	Gly	Ile	Leu	Ala	Lys	Ala
TAC	CTC	AAG	GGC	ATT	CTC	GCG	AAG	GCC
				130				
Pro	Phe	Arg	Val	Glu	Gly	Asp	Lys	Ala
CCG	TTC	CGC	GTC	GAA	GGC	GAC	AAG	GCC
					140			
Tyr	Gly	Pro	Gly	Ile	Ala	Asp	Asp	Lys
TAC	GGC	CCG	GGC	ATC	GCC	GAC	GAC	AAG
						150		
Gly	Gly	Asn	Ala	Val	Ile	Leu	His	Thr
GGC	GGC	AAC	GCG	GTC	ATC	CTG	CAC	ACG
							160	
Leu	Lys	Leu	Leu	Lys	Glu	Tyr	Gly	Val
CTC	AAG	CTG	CTG	AAG	GAA	TAC	GCC	GTC
								170
Arg	Asp	Tyr	Gly	Thr	Ile	Thr	Val	Leu
CGC	GAC	TAC	GGC	ACC	ATC	ACC	GTC	CTG
Phe	Asn	Thr	Asp	Glu	Glu	Lys	Gly	Ser
TTC	AAC	ACC	GAC	GAG	GAA	AAG	GGT	TCC
180								
Phe	Gly	Ser	Arg	Asp	Leu	Ile	Gln	Glu
TTC	GGC	TCG	CGC	GAC	CTG	ATC	CAG	GAA
	190							
Glu	Ala	Lys	Leu	Ala	Asp	Tyr	Val	Leu
GAA	GCC	AAG	CTG	GCC	GAC	TAC	GTC	CTC
		200						
Ser	Phe	Glu	Pro	Thr	Ser	Ala	Gly	Asp
TCC	TTC	GAG	CCC	ACC	AGC	GCA	GCC	GAC

Glu	Lys	Leu	210 Ser	Leu	Gly	Thr	Ser	Gly
GAA	AAA	CTC	TCG	CTG	GGC	ACC	TCG	GGC
Ile	Ala	Tyr	Val	220 Gln	Val	Asn	Ile	Thr
ATC	GCC	TAC	GTG	CAG	GTC	AAC	ATC	ACC
Gly	Lys	Ala	Ser	His	230 Ala	Gly	Ala	Ala
GGC	AAG	GCC	TCG	CAT	GCC	GGC	GCC	CCG
Pro	Glu	Leu	Gly	Val	Asn	240 Ala	Leu	Val
CCC	GAG	CTG	GGC	CTG	AAC	CCG	CTG	GTC
Glu	Ala	Ser	Asp	Leu	Val	Leu	250 Arg	Thr
GAG	GCT	TCC	GAC	CTC	GTG	CTG	CGC	ACG
Met	Asn	Ile	Asp	Asp	Lys	Ala	Lys	260 Asn
ATG	AAC	ATC	GAC	GAC	AAG	CCG	AAG	AAC
Leu	Arg	Phe	Asn	Trp	Thr	Ile	Ala	Lys
CTG	CGC	TTC	AAC	TGG	ACC	ATC	GCC	AAG
270 Ala	Gly	Asn	Val	Ser	Asn	Ile	Ile	Pro
GCC	GGC	AAC	GTC	TCG	AAC	ATC	ATC	CCC
Ala	280 Ser	Ala	Thr	Leu	Asn	Ala	Asp	Val
GCC	AGC	GCC	ACG	CTG	AAC	GCC	GAC	GTG
Arg	Tyr	290 Ala	Arg	Asn	Glu	Asp	Phe	Asp
CGC	TAC	CCG	CGC	AAC	GAG	GAC	TTC	GAC
Ala	Ala	Met	300 Lys	Thr	Leu	Glu	Glu	Arg
GCC	GCC	ATG	AAG	ACG	CTG	GAA	GAG	CGC
Ala	Gln	Gln	Lys	310 Lys	Leu	Pro	Glu	Ala
CCG	CAG	CAG	AAG	AAG	CTG	CCC	GAG	CCC

Asp	Val	Lys	Val	Ile	320 Val	Thr	Arg	Gly
GAC	GTG	AAG	GTG	ATC	GTC	ACG	CGC	GGC
Arg	Pro	Ala	Phe	Asn	Ala	330 Gly	Glu	Gly
CGC	CCG	GCC	TTC	AAT	GCC	GGC	GAA	GGC
Gly	Lys	Lys	Leu	Val	Asp	Lys	340 Ala	Val
GGC	AAG	AAG	CTG	GTC	GAC	AAG	CCG	GTG
Ala	Tyr	Tyr	Lys	Glu	Ala	Gly	Gly	350 Thr
GCC	TAC	TAC	AAG	GAA	GCC	GGC	GGC	ACG
Leu	Gly	Val	Glu	Glu	Arg	Thr	Gly	Gly
CTG	GGC	GTG	GAA	GAG	CGC	ACC	GGC	GGC
360 Gly	Thr	Asp	Ala	Ala	Tyr	Ala	Ala	Leu
GGC	ACC	GAC	GGC	GCC	TAC	GCC	GGC	CTC
Ser	370 Gly	Lys	Pro	Val	Ile	Glu	Ser	Leu
TCA	GGC	AAG	CCA	GTG	ATC	GAG	AGC	CTG
Gly	Leu	380 Pro	Gly	Phe	Gly	Tyr	His	Ser
GGC	CTG	CCG	GGC	TTC	GGC	TAC	CAC	AGC
Asp	Lys	Ala	390 Glu	Tyr	Val	Asp	Ile	Ser
GAC	AAG	GCC	GAG	TAC	GTG	GAC	ATC	AGC
Ala	Ile	Pro	Arg	400 Arg	Leu	Tyr	Met	Ala
GGC	ATT	CCG	CGC	CGC	CTG	TAC	ATG	GCT
Ala	Arg	Leu	Ile	Met	410 Asp	Leu	Gly	Ala
GGC	CGC	CTG	ATC	ATG	GAT	CTG	GGC	GCC
Gly	Lys							
GGC	AAG	TGA - 3'						

9. A recombinant DNA transfer vector according to any preceding claim characterised in that the transfer vector is a plasmid.
10. A recombinant DNA transfer vector according to claim 9  
5 whenever taken together with claim 7 having the designation pNM1, pNM111, pNM14, pNM21, pNM22, pNM31, pNM32 or pLEC3.
11. A microorganism transformed by a transfer vector characterised in that the transfer vector is a recombinant DNA transfer vector according to claim 1.
- 10 12. A microorganism according to claim 11 characterised in that the transfer vector is a recombinant DNA transfer vector according to claim 4.
13. A microorganism according to either claim 11 or claim 12 which is a bacterium of the species E.coli, Pseudomonas or  
15 Bacillus or a yeast of the species Saccharomyces cerevisiae.
14. A process for the preparation of a gene product characterised by
- (a) culturing a microorganism according to claim 12 in a culture medium to produce the gene product in the culture  
20 medium or the periplasmic space of the microorganism, and
- (b) isolating the gene product from the culture medium or the periplasmic space of the microorganism.
15. A process according to claim 14 characterised in that the gene product is Pseudomonas carboxypeptidase G<sub>2</sub> or E.coli  $\beta$  - galactosidase.


*Fig1.*

RESTRICTION ENZYME MAP OF pNM1

— REPRESENTS pBR322




2/7



RESTRICTION ENZYME MAP OF pNM111

*Fig.2.*


3/7



RESTRICTION ENZYME MAP OF pNM14


*Fig.3.*

4/7



RESTRICTION ENZYME MAP OF pNM21

*Fig.4.*



RESTRICTION ENZYME MAP OF pNM22

*Fig.5.*

6/7









Fig.6.



-  pBR 322
-  Yeast 2 $\mu$  plasmid
-  Yeast chromosomal leu 2 gene
-  Pseudomonas carboxypeptidase G2 gene

*Fig.7*

RESTRICTION ENZYME MAP OF pLEC 3



European Patent  
Office

# EUROPEAN SEARCH REPORT

01 21 352

Application number

DOCUMENTS CONSIDERED TO BE RELEVANT			EP 84301468.9
Category	Citation of document with indication, where appropriate, of relevant passages	Relevant to claim	CLASSIFICATION OF THE APPLICATION (Int. Cl. 7)
A, D	EP - A2 - 0 001 931 (GENENTECH, INC.) * Claims 1,6 *	1,5	C 12 N 15/00 C 12 P 19/34 C 07 C 103/52 C 12 P 21/00
A	EP - A2 - 0 049 619 (ELI LILLY AND COMPANY) * Claims 1,3 *	1,5	C 12 N 9/48 C 12 N 9/38// C 12 R 1/19 C 12 R 1/38 C 12 R 1/07 C 12 R 1/865
D, A	& GB-A-2 007 675 -----		
			TECHNICAL FIELDS SEARCHED (Int. Cl. 7)
			C 12 N C 12 P C 07 C 103/00
The present search report has been drawn up for all claims			
Place of search VIENNA		Date of completion of the search 02-08-1984	Examiner WOLF
<p><b>CATEGORY OF CITED DOCUMENTS</b></p> <p>X : particularly relevant if taken alone Y : particularly relevant if combined with another document of the same category A : technological background O : non-written disclosure P : intermediate document</p> <p>T : theory or principle underlying the invention E : earlier patent document, but published on, or after the filing date D : document cited in the application L : document cited for other reasons &amp; : member of the same patent family, corresponding document</p>			

(18)



Europäisches Patentamt  
European Patent Office  
Office européen des brevets

(11) Publication number:

**0 121 352  
B1**

(12)

## EUROPEAN PATENT SPECIFICATION

(45) Date of publication of patent specification: 07.01.88

(21) Application number: 84301468.9

(22) Date of filing: 06.03.84

(51) Int. Cl.<sup>4</sup>: C 12 N 15/00, C 12 N 9/38,  
C 12 N 9/52

(54) A leader sequence to promote the secretion of gene products.

(28) Priority: 28.03.83 GB 8308483

(43) Date of publication of application:  
10.10.84 Bulletin 84/41

(45) Publication of the grant of the patent:  
07.01.88 Bulletin 88/01

(84) Designated Contracting States:  
BE CH DE FR GB IT LI NL SE

(56) References cited:  
EP-A-0 001 931  
EP-A-0 049 619

(78) Proprietor: The Public Health Laboratory  
Service Board  
61 Colindale Avenue  
London NW9 5EQ (GB)

(72) Inventor: Atkinson, Anthony  
Twigley Mill Corner Winterbourne Gunner  
Salisbury Wiltshire SP 4 6JJ (GB)  
Inventor: Minton, Nigel Peter  
6 Highbury Avenue  
Salisbury Wiltshire SP2 7EX (GB)  
Inventor: Sherwood, Roger Franklin  
35 Attwood Road  
Salisbury Wiltshire SP1 3PR (GB)

(74) Representative: James, Stephen Richard et al  
Procurement Executive Ministry of Defence  
Patents 1A(4), Room 2014 Empress State  
Building  
Lillie Road London SW6 1TR (GB)

**EP 0 121 352 B1**

Note: Within nine months from the publication of the mention of the grant of the European patent, any person may give notice to the European Patent Office of opposition to the European patent granted. Notice of opposition shall be filed in a written reasoned statement. It shall not be deemed to have been filed until the opposition fee has been paid. (Art. 99(1) European patent convention).

Courier Press, Leamington Spa, England.



## Description

The present invention relates to fragments of specific deoxyribonucleotide sequences that promote the secretion of gene products from cells and in particular to recombinant DNA transfer vectors that contain these fragments.

Recent developments in biochemistry have led to the construction of recombinant DNA transfer vectors in which, transfer vectors, for example plasmids, are made to contain exogenous DNA. In some cases the recombinant incorporates heterologous DNA that codes for polypeptides that are ordinarily not produced by the organism susceptible to transformation by the recombinant vehicle.

In its basic outline a method of endowing a microorganism with the ability to synthesise a new protein involves three general steps:

(a) isolation and purification of the specific gene or nucleotide sequences containing the genetically coded information for the amino acid sequence of the desired protein or polypeptide,

(b) recombination of the isolated gene or nucleotide sequence with an appropriate transfer vector, typically DNA of a bacteriophage or plasmid to form a recombinant transfer vector that codes, in part, for the production of the desired protein or polypeptide,

(c) transfer of the vector to the appropriate microorganism and selection of a strain of the recipient microorganism containing the desired genetic information.

Provided the gene or nucleotide sequence expresses its protein or polypeptide in the chosen microorganism, growth of the microorganism should then produce the desired protein or polypeptide in significant quantities.

Once the microorganism has been cultured, the protein or polypeptide must be isolated from the undesired materials. This step is considerably facilitated if the majority of the desired protein or polypeptide is present in the culture medium and/or the periplasmic space of the microorganism. In other words purification may be performed in a more efficient manner if, once expressed, the protein or polypeptide passes through the cell membrane and out of the cytoplasm.

The passage of the protein or polypeptide through the cell membrane is desirable for two main reasons. First the desired protein or polypeptide will generally be foreign to the microorganism in which it is expressed. In many cases, therefore, it will be quickly broken down by proteolytic enzymes etc. in the cells cytoplasm and will, subsequently, have a short half life within the cell. By transferring the protein or polypeptide out of the cytoplasm soon after expression the stability of the protein or polypeptide will be greatly increased. Second the number of unwanted genetic materials and products (from which the desired protein or polypeptide must be isolated) will be far greater in the cell's cytoplasm than in the culture medium and/or in the cell's periplasmic space. It can be seen that on both of the above counts the transfer of the protein or polypeptide through the cell membrane and out of the cytoplasm will greatly facilitate protein or polypeptide isolation.

One way in which the secretion of gene products from the cell's cytoplasm may be promoted is to produce, within the cytoplasm, a preprotein or prepolyptide in which the desired protein or polypeptide is preceded by a signal polypeptide. The predominantly hydrophobic signal polypeptide directs the desired protein or polypeptide to the cell's periplasmic space, where the signal peptide is removed as the desired protein or polypeptide traverses the cell membrane.

Many of the known signal peptides contain cysteine residues. These residues have been found to react in the cell membrane and thereby inhibit the efficient transfer of the desired gene product out of the cell.

It is the primary object of the present invention to provide recombinant DNA transfer vectors containing a leader sequence polynucleotide that codes for a signal peptide that is cysteine free. Other objects and advantages of the present invention will become apparent from the following description thereof.

According to the present invention there is provided a recombinant DNA transfer vector comprising a leader sequence polynucleotide coding for signal polypeptide of formula I,

Met-Arg-Pro-Ser-Ile-His-Arg-Thr-Ala-Ile-Ala-  
Ala-Val-Leu-Ala-Thr-Ala-Phe-Val-Ala-Gly-Thr

The transfer vector may be a bacteriophage or, which is preferred, a plasmid.

Preferably the majority of the codons in the nucleotide sequence are those preferred for the expression of microbial genomes. Suitable codons are listed in UK 1,568,047 and UK 2007675A, and these publications are therefore incorporated herein by reference.

In one preferred embodiment of the present transfer vector the nucleotide sequence has formula II

5'	ATG	CGC	CCA	TCC	ATC	CAC	CGC	ACA	
	GCC	ATC	GCC	GCC	GTG	CTG	GCC	ACC	II
	GCC	TTC	GTG	GCG	GGC	ACC	— 3'		

The nucleotide sequence coding for the signal polypeptide (the leader sequence polynucleotide) will preferably be downstream of and in reading phase with a bacterial or a yeast promoter and a prokaryotic

ribosome binding site in the transfer vector. Moreover the leader sequence polynucleotide will either be upstream of an insertion site for a structural gene or, which is preferred, will be upstream of and in reading phase with a structural gene coding for a desired protein or polypeptide. Preferably the gene codes for a eukaryotic, particularly a mammalian, protein or polypeptide.

5 The structural gene may code for such prokaryotic proteins as *E. coli*  $\beta$ -galactosidase or *Pseudomonas* carboxy peptidase G<sub>2</sub> (CPG<sub>2</sub>) (Carboxypeptidase G<sub>2</sub> is an enzyme, produced by *Pseudomonas* species strain RS-16, that has application in cancer chemotherapy. It is a Zn<sup>2+</sup> containing dimer of 2×42,000 daltons and has high affinities (K<sub>m</sub> values of 10<sup>-5</sup> or 10<sup>-6</sup>M) for both 5-methyltetrahydrofolate, the predominant  
10 circulatory form of folate in mammals and for the folic acid antagonist methotrexate (MTX), which is widely used in cancer chemotherapy. The enzyme may be used directly for the plasma depletion of reduced folates, essential as co-factors in purine and particularly in pyrimidine biosynthesis. CPG<sub>2</sub> has been shown to inhibit the development of the Walker 256 carcinoma *in vivo* and to remove MTX from circulation in patients where prolonged exposure to high doses of MTX leads to toxicity).

15 Examples of transfer vectors according to the present invention that code for CPG<sub>2</sub> are pNM1, pNM111, pNM14, pNM21, pNM22, pNM31, pNM32 and pLEC3.

The promoter is preferably a high expression bacterial or yeast promoter for the structural gene in a variety of hosts. The particular choice of promoter will depend on the microorganism to be transformed. For example the transformation of *E. coli* will generally be effected by a transfer vector in which an *E. coli* promoter controls the expression of the structural gene. Examples of *E. coli* promoters are those present in  
20 the plasmids pBR 322 and pAT 153. By contrast, the transformation of *Pseudomonas* species will generally be effected by a transfer vector in which a *Pseudomonas* promoter controls the expression of the structural gene. Examples of *Pseudomonas* promoters are those present in the plasmid pKT 230 or *Pseudomonas* chromosomal DNA.

In order to express the structural gene the present transfer vector will be transformed into a suitable  
25 microorganism. According to a further aspect of the present invention therefore there is provided a microorganism transformed by a recombinant DNA transfer vector according to this invention. The microorganism will preferably be a bacterium or yeast in which high expression of the structural gene, within the transfer vector, occurs. Depending on the choice of promoter the microorganism may be a strain chosen from one of the following bacteria *E. coli*, *Pseudomonas* or the yeast *Saccharomyces cerevisiae*.

30 Having transformed the microorganism, the protein or polypeptide, for which the structural gene codes, may then be expressed by culturing the transformed microorganism in a culture medium. It is the primary advantage of the present invention that culturing the transformed microorganism affords a preprotein or prepolyptide in which the desired protein or polypeptide is preceded by the present signal polypeptide. This means that soon after expression the signal polypeptide directs the desired protein or  
35 polypeptide to the cell's periplasmic space, where the signal polypeptide is removed as the desired protein or polypeptide traverses the cell membrane. Since the present signal polypeptide is free of cysteine residues the desired gene product will be efficiently secreted through the membrane.

The present transfer vectors may be prepared by any of the methods that are well known in the recombinant DNA art. For example the leader sequence polynucleotide may be synthesised by the  
40 modified triester method of K. Itakura et al, *JACS*, 1975, 97, 7327 or by the improved oligodeoxynucleotide preparation described in UK 2007675A. The disclosure of both of these references is incorporated herein by reference. The synthesised polynucleotide may then be inserted in a transfer vector, preferably a plasmid. In the transfer vector it will preferably be downstream of and in reading phase with a bacterial or a yeast promoter and a prokaryotic ribosome binding site. The leader sequence polynucleotide should also be  
45 either upstream of a structural gene insertion site or upstream of and in reading phase with a structural gene.

Alternatively, DNA fragments containing the leader sequence polynucleotide may be obtained from natural sources, in particular from the chromosomal DNA of *Pseudomonas* species strain RS-16. In this particular case a polynucleotide (formula II above) coding for the present signal polypeptide immediately  
50 precedes a structural gene coding for CPG<sub>2</sub>. A number of the DNA fragments containing this leader sequence polynucleotide may therefore be recognised by their ability, on insertion into a plasmid and transformation of a microorganism by the resultant recombinant vector, to enable a microorganism to grow on folate. Examples of such recombinant transfer vectors that contain both a polynucleotide coding for the present signal polypeptide (formula II above) and a structural gene coding for CPG<sub>2</sub> are pNM1,  
55 pNM111, pNM14, pNM21, pNM22, pNM31, pNM32 and pLEC3. Of course, once a *Fol*<sup>+</sup> recombinant vector has been obtained in this way it may be subcloned to afford alternative vectors (either *Fol*<sup>+</sup> or *Fol*<sup>-</sup>) that also contain a polynucleotide coding for the present signal polypeptide.

Once a suitable DNA fragment has been isolated it may then be inserted in a transfer vector, preferably a plasmid. In the transfer vector the leader sequence polynucleotide on the inserted fragment should be  
60 downstream of and in reading phase with a bacterial or a yeast promoter and a prokaryotic ribosome binding site. The leader sequence polynucleotide should also be either upstream of a structural gene insertion site or upstream of and in reading phase with a structural gene.

The structural gene for insertion downstream of and in reading phase with the present leader sequence polynucleotide may be obtained, for example, by the synthetic methods mentioned above (this is  
65 particularly useful for the preparation of genes coding for small proteins). Alternatively the structural gene

# O 121 352

may be prepared from m=RNA by the use of the enzyme reverse transcriptase or may be isolated from natural sources (chromosomal DNA).

An example of the latter method is the isolation of DNA fragments containing a polynucleotide sequence (shown in Table 1) coding for the enzyme CPG<sub>2</sub> (amino acid sequence also shown in Table 1) from *Pseudomonas* species strain RS-16 chromosomal DNA. Examples of plasmids containing a CPG<sub>2</sub> structural gene, as well as a polynucleotide coding for the present signal polypeptide (formula II above), are pNM1, pNM111, pNM14, pNM21, pNM22, pNM31, pNM32 and pLEC3.

Once prepared or isolated the leader sequence polynucleotide and the structural gene will be inserted into a transfer vector, preferably a plasmid, to form a recombinant DNA transfer vector according to the present invention. The insertion step or steps will preferably be effected by one of the well known techniques in this art that employ restriction endonucleases, see for example the methods discussed in UK 2090600A, the disclosure of which is incorporated herein by reference. The choice of transfer vector will be determined by the microorganism in which the leader sequence polynucleotide and structural gene are to be expressed. Generally the transfer vector will be a cloning vehicle that is suitable for transforming the chosen microorganisms and that displays a phenotypical characteristic, such as antibiotic resistance, by which the recombinant transfer vectors may be selected. Thus, if the microorganism is to be *E. coli*, then suitable transfer vectors will be the *E. coli* plasmids pBR322 and pAT153. Alternatively, if the microorganism is to be *Pseudomonas* then a suitable transfer vector will be *Pseudomonas* pKT230.

TABLE 1  
A polynucleotide sequence, coding for CPG<sub>2</sub>, isolated from  
*Pseudomonas* species strain RS-16 chromosomal DNA

25	5'—	1 Met ATG	Arg CGC	Pro CCA	Ser TCC	Ile ATC	His CAC	Arg CGC	Thr ACA
30	Ala GCC	10 Ile ATCC	Ala GCC	Ala GCC	Val GTG	Leu CTG	Ala GCC	Thr ACC	Ala GCC
35	Phe TTC	20 Val GTG	Ala GCG	Gly GGC	Thr ACC	Ala GCC	Leu CTG	Ala GCC	Gln CAG
40	Lys AAG	30 Arg CGC	Asp GAC	Asn AAC	Val GTG	Leu CTG	Phe TTC	Gln CAG	Ala GCA
45	Ala GCT	40 Thr ACC	Asp GAC	Glu GAG	Gln CAG	Pro CCG	Ala GCC	Val GTG	Ile ATC
50	Lys AAG	50 Thr ACG	Leu CTG	Glu GAG	Lys AAG	Leu CTG	Val GTC	Asn AAC	Ile ATC
55	Glu GAG	60 Thr ACC	Gly GGC	Thr ACC	Gly GGT	Asp GAC	Ala GCC	Glu GAG	Gly GGC
60	Ile ATC	70 Ala GCC	Ala GCT	Ala GCG	Gly GGC	Asn AAC	Phe TTC	Leu CTC	Glu GAG
65	Ala GCC	80 Glu GAG	Leu CTC	Lys AAG	Asn AAC	Leu CTC	Gly GGC	Phe TTC	Thr ACG
	Val GTC	90 Thr ACG	Arg CGA	Ser AGC	Lys AAG	Ser TCG	Ala GCC	Gly GGC	Leu CTG
	Val GTG		Gly GGC	Asp GAC	Asn AAC	Ile ATC	Val GTG	Gly GGC	Lys AAG

# 0 121 352

TABLE 1 (contd.)

	Ile ATC	100 Lys AAG	Gly GGC	Arg CGC	Gly GGC	Gly GGC	Lys AAG	Asn AAC	Leu CTG
5									
	Leu CTG	Leu CTG	110 Met ATG	Ser TCG	His CAC	Met ATG	Asp GAC	Thr ACC	Val GTC
10									
	Tyr TAC	Leu CTC	Lys AAG	120 Gly GGC	Ile ATT	Leu CTC	Ala GCG	Lys AAG	Ala GCC
15									
	Pro CCG	Phe TTC	Arg CGC	130 Glu GAA	Gly GGC	Asp GAC	Lys AAG	Ala GCC	
20									
	Tyr TAC	Gly GGC	Pro CCG	Gly GGC	Ile ATC	140 Ala GCC	Asp GAC	Asp GAC	Lys AAG
25									
	Gly GGC	Gly GGC	Asn AAC	Ala GCG	Val GTC	Ile ATC	150 Leu CTG	His CAC	Thr ACG
30									
	Leu CTC	Lys AAG	Leu CTG	Leu CTG	Lys AAG	Glu GAA	Tyr TAC	160 Gly GGC	Val GTG
35									
	Arg CGC	Asp GAC	Tyr TAC	Gly GGC	Thr ACC	Ile ATC	Thr ACC	Val GTG	170 Leu CTG
40									
	Phe TTC	Asn AAC	Thr ACC	Asp GAC	Glu GAG	Glu GAA	Lys AAG	Gly GGT	Ser TCC
45									
	180 Phe TTC	Gly GGC	Ser TCG	Arg CGC	Asp GAC	Leu CTG	Ile ATC	Gln CAG	Glu GAA
50									
	Glu GAA	190 Ala GCC	Lys AAG	Leu CTG	Ala GCC	Asp GAC	Tyr TAC	Val CTG	Leu CTC
55									
	Ser TCC	Phe TTC	200 Glu GAG	Pro CCC	Thr ACC	Ser AGC	Ala GCA	Gly GGC	Asp GAC
60									
	Glu GAA	Lys AAA	Leu CTC	210 Ser TCG	Leu CTG	Gly GGC	Thr ACC	Ser TCG	Gly GGC
65									
	Ile ATC	Ala GCC	Tyr TAC	Val GTG	220 Gln CAG	Val GTC	Asn AAC	Ile ATC	Thr ACC
70									
	Gly GGC	Lys AAG	Ala GCC	Ser TCG	His CAT	230 Ala GCC	Gly GGC	Ala GCC	Ala GCC
75									
	Pro CCC	Glu GAG	Leu CTG	Gly GGC	Val GTG	Asn AAC	240 Ala GCG	Leu CTG	Val GTC

# 0 121 352

TABLE 1 (contd.)

	Glu GAG	Ala GCT	Ser TCC	Asp GAC	Leu CTC	Val GTG	Leu CTG	250 Arg CGC	Thr ACG
5									
	Met ATG	Asn AAC	Ile ATC	Asp GAC	Asp GAC	Lys AAG	Ala GCG	Lys AAG	260 Asn AAC
10	Leu CTG	Arg CGC	Phe TTC	Asn AAC	Trp TGG	Thr ACC	Ile ATC	Ala GCC	Lys AAG
	270 Ala GCC	Gly GGC	Asn AAC	Val GTC	Ser TCG	Asn AAC	Ile ATC	Ile ATC	Pro CCC
15									
	Ala GCC	280 Ser AGC	Ala GCC	Thr ACG	Leu CTG	Asn AAC	Ala GCC	Asp GAC	Val GTG
20									
	Arg CGC	Tyr TAC	290 Ala GCG	Arg CGC	Asn AAC	Glu GAG	Asp GAC	Phe TTC	Asp GAC
25									
	Ala GCC	Ala GCC	Met ATG	300 Lys AAG	Thr ACG	Leu CTG	Glu GAA	Glu GAG	Arg CGC
30									
	Ala GCG	Gln CAG	Gln CAG	Lys AAG	310 Lys AAG	Leu CTG	Pro CCC	Glu GAG	Ala GCC
35									
	Asp GAC	Val GTG	Lys AAG	Val GTG	Ile ATC	320 Val GTC	Thr ACG	Arg CGC	Gly GGC
40									
	Arg CGC	Pro CCG	Ala GCC	Phe TTC	Asn AAT	Ala GCC	330 Gly GGC	Glu GAA	Gly GGC
45									
	Gly GGC	Lys AAG	Lys AAG	Leu CTG	Val GTC	Asp GAC	Lys AAG	340 Ala GCG	Val GTG
50									
	Ala GCC	Tyr TAC	Tyr TAC	Lys AAG	Glu GAA	Ala GCC	Gly GGC	Gly GGC	350 Thr ACG
55									
	Leu CTG	Gly GGC	Val GTG	Glu GAA	Glu GAG	Arg CGC	Thr ACC	Gly GGC	Gly GGC
60									
	360 Gly GGC	Thr ACC	Asp GAC	Ala GCG	Ala GCC	Tyr TAC	Ala GCC	Ala GCG	Leu CTC
65									
	Ser TCA	370 Gly GGC	Lys AAG	Pro CCA	Val GTG	Ile ATC	Glu GAG	Ser AGC	Leu CTG
70									
	Gly GGC	Leu CTG	380 Pro CCG	Gly GGC	Phe TTC	Gly GGC	Tyr TAC	His CAC	Ser AGC

TABLE 1 (contd.)

	Asp	Lys	Ala	390 Glu	Tyr	Val	Asp	Ile	Ser
5	GAC	AAG	GCC	GAG	TAC	GTG	GAC	ATC	AGC
	Ala	Ile	Pro	400 Arg	Arg	Leu	Tyr	Met	Ala
	GCG	ATT	CCG	CGC	CGC	CTG	TAC	ATG	GCT
10	Ala	Arg	Leu	Ile	Met	410 Asp	Leu	Gly	Ala
	CGC	CGC	CTG	ATC	ATG	GAT	CTG	GGC	GCC
15	Gly	Lys	TGA	—3'					
	GGC	AAG							

Amino acids 1 to 22 are the present signal polypeptide.

Amino acids 23 to 415 are the CPG<sub>2</sub> structural gene.

NB The leader sequence polynucleotide is the preferred polynucleotide of formula II.

20 The present recombinant DNA transfer vectors, microorganisms transformed by the present recombinant DNA transfer vectors and processes for the preparation of said vectors and microorganisms will now be described by way of example only, with particular reference to the Figures in which:

Figure 1 is a restriction enzyme cleavage site map of pNM1,  
 25 Figure 2 is a restriction enzyme cleavage site map of pNM111,  
 Figure 3 is a restriction enzyme cleavage site map of pNM14,  
 Figure 4 is a restriction enzyme cleavage site map of pNM21,  
 Figure 5 is a restriction enzyme cleavage site map of pNM22, and  
 Figure 6 illustrates the process for the preparation of a recombinant plasmid containing both the  
 30 present leader sequence polynucleotide and the  $\beta$ -Galactosidase structural gene, and  
 Figure 7 is a restriction enzyme cleavage site map of pLEC3.

#### Materials and methods

##### Bacterial strains and plasmids

35 The bacterial strains used were *Escherichia coli* W5445 (*pro leu thi thr<sup>s</sup> sup E44 lac Y ton A r<sup>-</sup> m<sup>-</sup> Str<sup>r</sup>*) *Pseudomonas putida* 2440 (*r<sup>-</sup>*) and *Pseudomonas* sp strain RS-16. The plasmids employed were pBR322 (F. Bolivar et al Gene, 1977, 2, 95), pAT153 (A. J. Twigg et al, Nature, 1980, 283, 216) and pKT230 (M. Bagdasarjan et al, Gene 1981, 16, 237) and pROG5 (R. F. Sherwood et al, The Molecular Biology of Yeast, 1979 Cold Spring Harbor Publications).

##### Media and culture conditions

40 *E. coli* was routinely cultured in L-broth (1% tryptone, 0.5% yeast extract, 0.5% NaCl). Solidified medium (L-agar) consisted of L-broth with the addition of 2% (w/v) agar (Bacto-Difco). Antibiotic concentrations used for the selection of transformants were 50  $\mu$ g/ml ampicillin, 15  $\mu$ g/ml tetracycline and  
 45 30  $\mu$ g/ml kanamycin. In the case of *E. coli* these were conducted in 2YT liquid medium (1.6% tryptone, 1% yeast extract, 0.5% NaCl) containing 1% glucose, and 0.05% folate where appropriate. The pseudomonads were grown in a minimal salts solution consisting of per litre: MgSO<sub>4</sub>, 0.05 g; CaCl<sub>2</sub>, 2H<sub>2</sub>O, 0.05 g; FeSO<sub>4</sub> · 7H<sub>2</sub>O, 0.005 g; MnSO<sub>4</sub>, 0.0015 g; Na<sub>2</sub>MbO<sub>4</sub>, 2H<sub>2</sub>O, 0.0015 g; KH<sub>2</sub>PO<sub>4</sub>, 5 g; K<sub>2</sub>HPO<sub>4</sub> · 3H<sub>2</sub>O, 12 g; glutamate, 10 g. The minimal medium employed for *E. coli* was M9 medium (J. Miller, Experiments in  
 50 molecular genetics, Cold Spring Harbor Laboratory, Cold Spring Harbor, New York, 1972).

##### Purification of DNA

Plasmids were purified from chloramphenicol amplified cultures (D. B. Clewell, J. Bacteriol, 1972, 110, 667) by Brij-lysis (D. B. Clewell et al, Proc. Natl. Acad. Sci., USA, 1969, 62, 1159) and subsequent caesium  
 55 chloride-ethidium bromide density gradient centrifugation (A. Colman et al, Eur. J. Biochem, 1978, 91, 303). A rapid, small scale plasmid isolation technique (Birnbolm et al. Nucl. Acids Res. 1979, 7, 1513) was also employed for screening purposes. Chromosomal DNA from the donor *Pseudomonas* strain (RS-16) was prepared essentially as described by J. Marmar, J. Mol. Biol, 1961, 3, 208.

##### Restriction, ligation and transformation methods

60 Restriction endonucleases and DNA ligase were purchased from Bethesda Research Laboratories and used in the buffers and under the conditions recommended by the supplier. Transformation of *E. coli* was essentially as described by S. N. Cohen et al., Proc. Natl. Acad. Sci., USA, 1972, 69, 2110, while *Ps. putida* was transformed by the method of M. Bagdasarjan and K. N. Timmis, Current Topics in Microbiology and  
 65 Immunology, Eds. P. H. Hofschneider and W. Goebel, Springer Verlag, Berlin, 1981, p. 47.

#### Agarose gel electrophoresis

Digests were electrophoresed in 0.8% agarose slab gels (10 cm×20 cm×0.5 cm) on a standard vertical system (Raven), employing Tris-borate-EDTA buffer. Electrophoresis of undigested DNA was at 125V, 50 mA for 3 hours, while digested DNA was electrophoresed at 15V, 10 mA for 16 hours. Fragment sizes were estimated by comparison with fragments of  $\lambda$ DNA digested with *Hind*III and  $\lambda$ DNA cut with both *Hind*III and *Eco*RI. Fragments were isolated from gels using electroelution (M. W. McDonnell et al, Proc. Natl. Acad. Sci, USA, 1977, 74, 4835).

#### Determination of carboxypeptidase G<sub>2</sub> activity

Bacteria were grown in 1 litre batch culture and 100 ml samples taken at various stages in the growth phase. Samples were cooled on ice, centrifuged at 13,000×g for 10 minutes and resuspended and frozen in 5 ml of 0.1 M Tris HCl, pH 7.3 containing 0.2 mM ZnSO<sub>4</sub>. The cells were disrupted using a MSE Ultrasonic Disintegrator (150 W) at medium frequency, amplitude 2, for three 30-second intervals on ice. Cell debris was removed by centrifugation at 10,000×g for 5 minutes. CPG<sub>2</sub> activity was determined after J. L. McCullough et al, J. Biol. Chem, 1971, 246, 7207. A 1 ml reaction cuvette containing 0.9 ml of 0.1 M Tris-HCl, pH 7.3 plus 0.2 mM ZnSO<sub>4</sub> and 0.1 ml of 0.6 mM methotrexate was equilibrated at 37°C. Enzyme extract was added to the test cuvette and the decrease in absorbance at 320 nm measured using a Pye-Unicam SP1800 double-beam spectrophotometer. Enzyme activity per ml extract was calculated as  $\Delta$  320 nm absorbance/min divided by 8.3, which is equivalent to the hydrolysis of 1  $\mu$ mol of MTX/min at 37°C. Protein concentration was determined by the method of M. M. Bradford, Anal. Biochem., 1976, 72, 248.

#### Cell fractionation techniques

Bacterial cultures were grown in the low phosphate medium of H. C. Neu and L. A. Heppel, (J. Biol. Chem., 1964, 240, 3685), supplemented with 100  $\mu$ g/ml ampicillin, to an OD<sub>450</sub>=1.0. 40 ml of culture was centrifuged at 5000 g for 10 min, washed in 5 ml of 10 mM Tris-HCl pH 7.0, and resuspended in 0.9 ml of 0.58 M sucrose, 0.2 mM DTT, 30 mM Tris-HCl pH 8.0. Conversion to spheroplasts was achieved by the addition of 20  $\mu$ l of lysozyme (2 mg/ml), 40  $\mu$ l 0.1 M EDTA, and incubation at 23° for 10 min (H. C. Neu et al, J. Biol. Chem., 1964, 239, 3893). The spheroplasts were placed on ice and 0.1 ml of 30% (w/v) BSA added, followed by 5 ml of sucrose-tris buffer. Sedimentation of the spheroplasts was achieved by centrifugation at 5000 g for 10 min and the supernatant retained as the 'periplasmic' fraction. The pellet was resuspended in 5 ml-10 mM Tris-HCl, 0.2 mM DTT pH 7.0 and sonicated at 20 Kc/sec, 2 Amps for 15 sec. Remaining whole cells were removed by centrifugation at 1000×g for 10 min. Centrifugation at 100000×g for 1 hr, at 4°C, separated the soluble (cytoplasmic) proteins from the particulate (membrane-bound) proteins. The membrane pellet was resuspended in 1 ml of 10 mM Tris-HCl, 0.2 mM DTT, pH 7.0.

CPG<sub>2</sub> was assayed as described. Alkaline phosphatase was assayed according to J. Miller, Experiments in Molecular genetics, Cold Spring Harbor Laboratory, Cold Spring Harbor, New York, 1972, NADH oxidase according to M. J. Osborn et al, J. Biol. Chem., 1972, 247, 3962 and glyceraldehyde-3-phosphate dehydrogenase after K. Suzuki et al, FEMS, 1971, 13, 217.

#### Example 1

Preparation of recombinant plasmid pNM1 (A plasmid containing both the present leader sequence polynucleotide and the CPG<sub>2</sub> structural gene).

To isolate the gene for carboxypeptidase G<sub>2</sub> together with the leader sequence polynucleotide chromosomal DNA prepared from the *Pseudomonas* host (strain RS-16) was partially digested with *Sau*3A and fragments of between 6—8 Md isolated from agarose gels by electroelution. The 'sized' DNA was ligated with alkaline phosphatase treated *Bam*HI cut pBR322, transformed into *E. coli* W5445, and Ap<sup>r</sup> transformants selected. Of the 3,500 Ap<sup>r</sup> colonies obtained, approximately 70% were Tc<sup>r</sup>. Utilisation of a rapid plasmid isolation technique on 50 Ap<sup>r</sup> Tc<sup>r</sup> transformants demonstrated that 90% of the gene bank harboured plasmids of the expected size. As a further check on the authenticity of the gene bank, the individual clones were screened for the acquisition of a Leu<sup>+</sup> phenotype. Two such clones were identified. Both carried a plasmid capable of transforming *leuB* (B-isopropylmalate dehydrogenase) *E. coli* mutants to prototrophy.

Acquisition of a functional CPG<sub>2</sub> gene should enable *E. coli* to utilise folic acid as a carbon source. The 2,400 gene bank clones were screened for the ability to grow on minimal medium containing folate as the sole source of carbon (i.e. Fol<sup>+</sup>). A single Fol<sup>+</sup> clone was detected and shown to harbour a plasmid capable of transforming plasmid-minus W5445 to the Fol<sup>+</sup> phenotype. Classical restriction mapping of this plasmid (pNM1) was undertaken which revealed the presence of a 5.9 Md insert of pseudomonad DNA within pBR322. The restriction enzyme cleavage site map of pNM1 is given in Figure 1. The nucleotide sequence of the leader sequence polynucleotide and the CPG<sub>2</sub> structural gene is given in Table 1.

#### Example 2

Subcloning of -plasmid pNM1 to form pNM111

In order to pinpoint the position of the CPG<sub>2</sub> gene and the leader sequence polynucleotide within the 5.9 Md insert, subcloning of various restriction enzyme fragments, into pBR322, was undertaken. A functional CPG<sub>2</sub> gene was shown not to occur on *Xho*I or *Sph*I fragments of the pNM1 insert, but was

present on a 3.1 Md *Bgl*II fragment. This latter fragment was cloned into the *Bam*HI site of pBR322 to give pNM11 (6.0 Md). A further reduction in the size of pNM11 was achieved by digesting with *Sa*I and relegating the resultant fragment to yield pNM111. In addition, plasmids in which the smaller 0.95 Md *Sa*I fragment had become inserted in the opposite orientation to the parent plasmid (pNM11) were *Fol*<sup>-</sup>. Taken together these subcloning results indicate that the CPG<sub>2</sub> gene and the leader sequence polynucleotide lie between the *Bgl*II site at 4.14 and the *SA*I site at 6.03 on Pnml. Furthermore, the gene contains a *Sph*I (5.17), *Sa*II (5.07) and at least one *Xho*I (4.56 and/or 5.56) site. The restriction enzyme cleavage site map of pNM111 is given in Figure 2.

#### 10 Example 3

Preparation of recombinant plasmid pNM14. (A plasmid containing both the present leader sequence polynucleotide and the CPG<sub>2</sub> structural gene)

The 3.1 Md *Bgl*II fragment from Example 2 above was partially digested with *Sau*3A. These fragments were then cloned into the *Bam*HI site of pAT153 and transformed into *E. coli* W5445. Of the two *Ap*<sup>r</sup> *Tc*<sup>s</sup> *Fol*<sup>-</sup> colonies obtained, one contained a plasmid which had acquired an extra *Sa*I and *Bam*HI site, this was pNM14. The restriction enzyme cleavage site map of pNM14 is given in Figure 3. Sequencing of the leader sequence polynucleotide and the CPG<sub>2</sub> structural gene present in pNM14 gave the nucleotide structure shown in Table 1. DNA sequencing of pNM14 also revealed that the *Sa*I—*Bam*HI fragment was a duplication of a segment of DNA from within the insert (marked \* on Figure 3) composed of two contiguous *Sau* 3A fragments.

#### Example 4 and 5

Preparation of recombinant plasmids pNM21 and pNM22 (Plasmids containing both the present leader sequence polynucleotide and the CPG<sub>2</sub> structural gene)

The 3.1 Md *Bgl*II fragment from Example 2 was cloned into the *Bam*HI site of pAT 153 and transformed into *E. coli* W5445. Two *Ap*<sup>r</sup> *Tc*<sup>s</sup> *Fol*<sup>-</sup> colonies were obtained, one containing a plasmid pNM21 in which the fragment was inserted in the opposite orientation to pNM1 and one containing a plasmid pNM22 in which the fragment was inserted in the same orientation as pNM1. The restriction enzyme cleavage site maps of pNM21 and pNM22 are given in Figures 4 and 5 respectively.

The two plasmids, pNM21 and pNM22 both transformed *E. coli* to *Fol*<sup>+</sup>, indicating that a pseudomonad promoter was present on the 3.1 Md fragment. However, cells carrying the plasmid pNM21, in which the *Bgl*II fragment was cloned in the opposite orientation to pNM1, exhibited more rapid growth with folic acid as the sole carbon source. This difference was clearly visible on agar medium, where colonies developed concentric yellow "halos" of precipitated pteric acid, the insoluble product of folate hydrolysis.

Confirmation that pNM21 gave enhanced expression of CPG<sub>2</sub> over pNM22, was obtained by assaying enzyme production during batch growth of cells containing either plasmid. (The cells were grown in complex medium supplemented with 1% (w/v) glucose and where appropriate 0.05% (w/v) folic acid. The generation time was 56—66 min. The culture was sampled at hourly intervals and whole cells were disrupted by sonication. Enzyme activity was determined in the centrifugal extract). Results are given in Table 2.

The expression of CPG<sub>2</sub> from the plasmids pNM22 and pNM1 was 2.5 units/litre of culture, representing 0.005% soluble protein. In contrast, expression from pNM21 was 3000—3500 units/litre of culture, which represented 4.7% soluble protein. As the cloned gene is inserted into the *Bam*HI site of pAT153, the observed higher expression of pNM21 is almost certainly due to transcriptional read through from the *Tc* promoter. The low expression of CPG<sub>2</sub> carried on plasmids pNM1 and pNM22 is consistent with the view that *Pseudomonas* promoters function poorly in *E. coli*. It is also apparent from Table 2 that in the presence of folate there is a two-fold increase in the specific activity of enzyme measured in cell sonicates. This phenomenon has been observed in all experiments, but does not seem to be associated with classical induction of the CPG<sub>2</sub> gene, as overall enzyme yield in the presence or absence of folate remains at about 3000 u/litre culture. It in fact reflects a consistent depression in the soluble protein levels measured in sonicates from cells grown in the presence of folate. There is no obvious difference in growth rate of cells grown with folate and the reasons for this result are not clear.



TABLE 2

Carboxypeptidase G production by *E. coli* W5445, containing the plasmids pNM1, pNM21 and pNM22

5	Culture	Carboxypeptidase G <sub>2</sub> specific activity (U/MG soluble protein)					
		pNM1		pNM22		pNM21	
	Age (hr)	-Fol	+Fol	-Fol	+Fol	-Fol	+Fol
10	1	—	—	—	—	11.5	13.4
	2	—	—	—	—	12.9	9.6
15	3	.008	.005	.010	.019	13.9	23.3
	4	.009	.011	.015	.013	12.3	26.9
	5	.007	.019	.016	.016	11.5	25.6
20	6	.005	.024	.014	.023	13.7	24.1
	7	.015	.029	.024	.043	13.2	20.6
25	8	.013	.028	.024	.046	13.0	23.6

Expression of the cloned gene in *Ps. putida*

The observation that the CPG<sub>2</sub> gene was expressed in *E. coli* regardless of the orientation of the gene within the vector suggested that the promoter region of the CPG<sub>2</sub> gene had been cloned with the structural gene and the leader sequence polynucleotide. The low expression of CPG<sub>2</sub> within *E. coli* from its natural promoter (pNM1, pNM22, pNM111) confirmed other findings that *Pseudomonas* promoters are poorly recognised by *E. coli* RNA polymerases. It would be expected that if the gene was introduced back into a pseudomonad cellular environment, then improved expression from the *Pseudomonas* promoter should result. The 3.1 Md *Bgl*III fragment was subcloned into the *Pseudomonas* cloning vector pKT230 at its single *Bam*HI site. Two plasmids were obtained, pNM31 and pNM32 representing the two possible orientations of the cloned gene. These plasmids were transformed into *Ps. putida* 2440 by the method of Bagdasarian and Timmis. *Pseudomonas* cells carrying both plasmids were cultured in minimal salts medium and enzyme production monitored.

Yields of 500–1000 units/litre of culture were obtained regardless of gene orientation within the plasmid. Specific activity of the enzyme in cell sonicates was 1.5 to 4.0 U/mg protein representing 0.3 to 0.7% soluble protein (compared with <0.05% soluble protein in the donor strain RS-16). This result strongly indicates that the CPG<sub>2</sub> promoter is present and operated in a pseudomonad background. When the same plasmids were transformed into *E. coli* W5445 12–40 Units/litre were found at specific activity <0.07 U/mg (<0.01% soluble protein).

Periplasmic localisation of CPG<sub>2</sub>

There is evidence that CPG<sub>2</sub> is located in or near the periplasmic space of *Pseudomonas* strain RS-16. Pteric acid, the product of CPG<sub>2</sub> hydrolysis of folic acid is extremely insoluble and is found predominantly outside the cell in both liquid and solid media. Exogenous pteric acid is also seen in *E. coli* cultures containing the cloned gene when folic acid is present in the medium. This is clearly demonstrated by the "halo" of precipitated pteric acid observed around colonies carrying plasmids in which expression of CPG<sub>2</sub> is from the Tc promoter of pBR322 (e.g. pNM21).

The localisation of CPG<sub>2</sub> produced by *E. coli* cells carrying pNM21 was examined by the separation of cellular proteins into cytoplasmic, periplasmic, and whole membrane fractions. As a control, levels of three marker enzymes, alkaline phosphatase (periplasmic), glyceraldehyde-3-phosphate dehydrogenase (cytoplasmic) and NADH · O<sub>2</sub> oxidoreductase (membrane-bound), were also determined. As can be seen from Table 3 97% of the CPG<sub>2</sub> activity occurs in the periplasm, equivalent to the marker periplasmic enzyme, alkaline phosphatase. This confirms the presence in pNM21 of a leader sequence polynucleotide next to the CPG<sub>2</sub> gene that codes for a signal polypeptide according to this invention that promotes the secretion of CPG<sub>2</sub> from the cytoplasm into the periplasmic space.

Carboxypeptidase G<sub>2</sub> synthesised in *E. coli*

The specific activity of CPG<sub>2</sub> in crude cell extracts of cells carrying pNM21 was 50-fold higher than equivalent extracts from *Pseudomonas* strain RS-16. To determine whether the cloned gene=product in *E.*

*coli* had the same properties as CPG<sub>2</sub> from the pseudomonad, enzyme was purified from *E. coli* carrying pNM21. The specific activity of purified CPG<sub>2</sub> (single band SDS-PAGE) was 535 U/mg of protein, which compares to 550 U/mg of protein from the pseudomonad. CPG<sub>2</sub> purified from *E. coli* clone pNM21 co-chromatographed with CPG<sub>2</sub> from *Pseudomonas* strain RS-16 at a sub-unit molecular weight value of 42,000 daltons. Km values using methotrexate as substrate were  $7.4 \times 10^{-6}$  M and  $8.0 \times 10^{-6}$  M respectively. In addition, antiserum raised against the *Pseudomonas* enzyme indicated immunological identity between the *E. coli* and *Pseudomonas* CPG<sub>2</sub>, as a confluent precipitation line was formed on Ouchterlony double diffusion analysis.

TABLE 3  
Localisation of carboxypeptidase

Fraction	CPG <sub>2</sub>	Enzyme activity		
		AP	GAPDH	NADHOX
Periplasmic	97.0	97.1	6.8	0.25
Cytoplasmic	2.6	2.3	93	8.4
Membrane-bound	0.4	0.6	0.2	89.1

AP=Alkaline phosphatase

GAPDH=Glyceraldehyde-3-phosphate dehydrogenase

NADHOX=NADH · O<sub>2</sub> oxidoreductase

#### Example 6

Preparation of a recombinant plasmid containing both the present leader sequence polynucleotide and the β-Galactosidase structural gene

Plasmid pNM14 (Example 3) was treated with *Sau* 3A (GATC) and the fragments were cloned into the *Bam* HI site of M13 mp7 template DNA (single stranded DNA (Step A of Figure 6). The product carrying a 318bp *Sau* 3A fragment coding for the present signal polypeptide and the first 22 amino acids of CPG<sub>2</sub> (nucleotide sequence of this fragment shown in Table 4) was selected and made double stranded. The DNA coding for the signal polypeptide (and the first part of CPG<sub>2</sub>) was then excised as an *Eco*RI fragment. This *Eco*RI fragment was then cloned into the promoter cloning vector *E. coli* pMC1403 (M. J. Casadaban *et al*, J. Bacteriol, 1980, 143, 971), which carries only the structural gene (*lac* Z) for β-galactosidase (i.e. no promoter and no ATG start codon) (Steps B and C of Figure 6). Plasmids were obtained in which the *Eco*RI fragment had inserted in both orientations, however, only those in which fusion of the CPG<sub>2</sub> sequence to the β-galactosidase sequence had occurred (i) yielded a 0.34 Kb fragment upon digestion with *Bam*HI; (ii) enabled the host cell to hydrolyse the colourless lactose analogue, BCIg, and impart a blue colouration to colonies. The 0.34 Kb *Bam*HI fragment has been recloned into M13mp7 and sequenced to confirm that fusion has occurred. The 'precursor' fusion produced will consist of the signal peptide, the first 22 amino acids of CPG<sub>2</sub>, 6 amino acids derived from the M13mp7 and pMC1403 linker units, and β-galactosidase from its 8th amino acid onward.

Localisation experiments have been performed on cells carrying a plasmid coding for the 'fusion gene' where the cellular proteins have been fractionated into periplasmic, cytoplasmic and membrane fractions. In these experiments an organism (*E. coli* MC 1061) which is deleted for the *lac* Z gene was grown in phosphate medium (H. C. Neu *et al*, J. Biol. Chem., 1964, 240, 3685) and periplasmic enzymes were released from the harvested cells by conversion to spheroplasts. Separation of soluble proteins (cytoplasmic) from particulate proteins (membrane band) was achieved by sonicating the harvested spheroplasts and subsequent centrifugation at 100,000 g for 1 hr, to sediment the cell membrane (T. J. Silhary *et al*, Proc. Natl. Acad. Sci. USA, 1976, 73, 3423).

The results given in Table 5 demonstrate the presence of 50% of the β-galactosidase activity in the periplasmic space. This result is in direct contrast to similar work involving fusion of other periplasmic protein signal sequences to β-galactosidase, where the fusion proteins are not exported, but become jammed in the membrane (P. J. Bassford *et al*, J. Bacteriol, 1979, 139, 19 and S. D. Emr *et al*, J. Cell. Biol., 1980, 86, 701).

# 0 121 352

TABLE 4  
The polynucleotide sequence of the 318 bp *Sau* 3A fragment from recombinant plasmid pNM14

5	5'—G	ATC	CAC	GCA	CTG	AAG	GCG	CGC	GGC
	AAG	ACG	CGC	GGC	GTG	GCG	ACG	CTG	TGC
	ATC	GGC	GGG	GGC	GAA	GGC	ACC	GCA	GTG
10	GCA	CTC	GAT	TGC	TAT	AAG	AAC	CAT	GGC
	TGG	GGA	CGC	CCG	ACA	ACA	GGC	GTC	CAC
15	CAG	CTT	TTT	TCA	TTC	CGA	CAA	CCC	GAA
	CGA	ACA	ATG	CGT	AGA	GCA	GGA	GAT	TCC
20		Met	Arg	Pro	Ser	Ile	His	Arg	Thr
		ATG	CGC	CCA	TCC	ATC	CAC	CGC	ACA
	Ala	Ile	Ala	Ala	Val	Leu	Ala	Thr	Ala
	GCC	ATC	GCC	GCC	GTG	CTG	GCC	ACC	GCC
25	Phe	Val	Ala	Gly	Thr	Ala	Leu	Ala	Gln
	TTC	GTG	GCG	GGC	ACC	GCC	CTG	GCC	CAG
	Lys	Arg	Asp	Asn	Val	Leu	Phe	Gln	Ala
	AAG	CGC	GAC	AAC	GTG	CTG	TTC	CAG	GCA
30	Ala	Thr	Asp	Glu	Gln	Pro	Ala	Val	Ile
	GCT	ACC	GAC	GAG	CAG	CCG	GCC	GTG	ATC

NB. This fragment carries the leader sequence coding for the signal polypeptide, a part of the CPG<sub>2</sub> structural gene coding for the first 22 amino acids of the protein, the ATG start codon, the CPG<sub>2</sub> ribosome binding site (AGGA) and other components of the CPG<sub>2</sub> promoter region.

TABLE 5  
Localisation of signal peptide— $\beta$ -galactosidase fusion protein

40		CPG <sub>2</sub> / $\beta$ -gal	% Localisation <sup>a</sup>		
			AP	GAPDH	NADHOX
45	Periplasmic	50.3	97.3	3.4	0.4
	Cytoplasmic	30.9	2.5	95.3	8.2
	Membrane-bound	18.8	0.2	1.3	89.4

<sup>a</sup>=average results from 4 experiments

CPG<sub>2</sub>/ $\beta$ -gal=Carboxypeptidase G<sub>2</sub>- $\beta$ -galactosidase fusion protein

AP=Alkaline phosphatase

GAPDH=Glyceraldehyde-3-phosphate dehydrogenase

NADHOX=NADH · O<sub>2</sub> Oxidoreductase

## Example 7

Preparation of a recombinant plasmid, containing both the present leader sequence polynucleotide and the CPG<sub>2</sub> structural gene, able to replicate in *E. coli* and *S. cerevisiae*

60 A 2.03 kilobase *Bam*HI fragment coding for the present signal polypeptide and the entire CPG<sub>2</sub> molecule was cloned in both orientations into the *Bam*HI site of an *E. coli*/*S. cerevisiae* shuttle vector pROG5 (R. F. Sherwood and R. K. Gibson, The Molecular Biology of Yeast, 1979, Cold Spring Harbor Publications) to give plasmids pLEC3 and pLEC4 (Figure 7). These plasmids were transformed into *S. cerevisiae* strain LL20 by the lithium acetate induced transformation method described by Ito *et al.*, J. Bact., 1983, 153, 163. Yields equivalent to 10–20 units/litre of culture volume were obtained regardless of gene

orientation within the plasmid. Specific activity of the enzyme in total cell extracts was 0.2—0.3 u/mg protein representing 0.005% soluble protein. This level of expression from the pseudomonad promoter in a yeast background is similar to the level found when the gene was reading from its own promoter in *E. coli* (0.01% soluble protein).

5 Localisation experiments have been performed on yeast cells carrying the above plasmids by sphaeroplasting the cells using standard techniques described by J. B. D. Beggs, Nature, 1978, 275, 105. Periplasmic enzymes, localised outside of the cell membrane, were released when the cell wall was removed. The osmotic stabiliser (1.2M sorbitol) was then replaced by 0.1M Tris-HCl buffer, pH 7.3 containing 0.2 mM ZnCl<sub>2</sub> to lyse the sphaeroplasts and the whole centrifuged at 100,000×g for 1 hour to  
10 separate proteins in the soluble cytoplasmic fraction from membrane bound proteins. The results in Table 6 demonstrate the presence of 64% of the CPG<sub>2</sub> activity in the periplasmic fraction and a further 16% associated with the cell membrane fraction.

TABLE 6  
Localisation of CPG<sub>2</sub> in *S. cerevisiae*

	% CPG <sub>2</sub> activity
Periplasmic	64
20 Cytoplasmic	20
Membrane bound	16

## 25 Claims

1. A recombinant DNA transfer vector comprising a leader sequence polynucleotide downstream of and in reading phase with a bacterial or yeast promoter and a prokaryotic ribosome binding site and upstream of and in reading phase with a structural gene, characterised in that the leader sequence  
30 polynucleotide codes for a signal polypeptide of formula I

Met-Arg-Pro-Ser-Ile-His-Arg-Thr-  
Ala-Ile-Ala-Ala-Val-Leu-Ala-Thr-  
Ala-Phe-Val-Ala-Gly-Thr

35

2. A recombinant DNA transfer vector according to claim 1 characterised in that the leader sequence polynucleotide is of formula II

5'—	ATG	CGC	CCA	TCC	ATC	CAC	CGC	ACA	
40	GCC	ATC	GCC	GCC	GTG	CTG	GCC	ACC	II
	GCC	TTC	GTG	GCG	GGC	ACC	—3'.		

3. A recombinant DNA transfer vector according to either claim 1 or claim 2 characterised in that the structural gene codes for Pseudomonas carboxypeptidase G<sub>2</sub> (CPG<sub>2</sub>).

45 4. A recombinant DNA transfer vector according to either claim 1 or claim 2 characterised in that the structural gene codes for a protein or polypeptide other than Pseudomonas carboxypeptidase G<sub>2</sub>.

5. A recombinant DNA transfer vector according to claim 4 characterised in that the structural gene codes for a prokaryotic protein other than Pseudomonas carboxypeptidase G<sub>2</sub>.

6. A recombinant DNA transfer vector according to claim 5 characterised in that the structural gene  
50 codes for *E. Coli* β-galactosidase.

7. A recombinant DNA transfer vector according to claim 3 comprising a polynucleotide of formula

55	5'—	1 Met ATG	Arg CGC	Pro CCA	Ser TCC	Ile ATC	His CAC	Arg CGC	Thr ACA
		10 Ala GCC	Ile ATC	Ala GCC	Ala GCC	Val GTG	Leu CTG	Ala GCC	Thr ACC
60			20 Phe TTC	Val GTG	Ala GCG	Gly GGC	Thr ACC	Ala GCC	Leu CTG
									Gln CAG

65

0 121 352

	Lys AAG	Arg CGC	Asp GAC	30 Asn AAC	Val GTG	Leu CTG	Phe TTC	Gln CAG	Ala GCA
5	Ala GCT	Thr ACC	Asp GAC	Glu GAG	40 Gln CAG	Pro CCG	Ala GCC	Val GTG	Ile ATC
10	Lys AAG	Thr ACG	Leu CTG	Glu GAG	Lys AAG	50 Leu CTG	Val GTC	Asn AAC	Ile ATC
15	Glu GAG	Thr ACC	Gly GGC	Thr ACC	Gly GGT	Asp GAC	60 Ala GCC	Glu GAG	Gly GGC
20	Ile ATC	Ala GCC	Ala GCT	Ala GCG	Gly GGC	Asn AAC	Phe TTC	70 Leu CTC	Glu GAG
	Ala GCC	Glu GAG	Leu CTC	Lys AAG	Asn AAC	Leu CTC	Gly GGC	Phe TTC	80 Thr ACG
25	Val GTC	Thr ACG	Arg CGA	Ser AGC	Lys AAG	Ser TCG	Ala GCC	Gly GGC	Leu CTG
30	90 Val GTG	Val GTG	Gly GGC	Asp GAC	Asn AAC	Ile ATC	Val GTG	Gly GGC	Lys AAG
35	Ile ATC	100 Lys AAG	Gly GGC	Arg CGC	Gly GGC	Gly GGC	Lys AAG	Asn AAC	Leu CTG
	Leu CTG	Leu CTG	110 Met ATG	Ser TCG	His CAC	Met ATG	Asp GAC	Thr ACC	Val GTC
40	Tyr TAC	Leu CTC	Lys AAG	120 Gly GGC	Ile ATT	Leu CTC	Ala GCG	Lys AAG	Ala GCC
45	Pro CCG	Phe TTC	Arg CGC	Val GTC	130 Glu GAA	Gly GGC	Asp GAC	Lys AAG	Ala GCC
50	Tyr TAC	Gly GGC	Pro CCG	Gly GGC	Ile ATC	140 Ala GCC	Asp GAC	Asp GAC	Lys AAG
55	Gly GGC	Gly GGC	Asn AAC	Ala GCG	Val GTC	Ile ATC	150 Leu CTG	His CAC	Thr ACG
	Leu CTC	Lys AAG	Leu CTG	Leu CTG	Lys AAG	Glu GAA	Tyr TAC	160 Gly GGC	Val GTG
60	Arg CGC	Asp GAC	Tyr TAC	Gly GGC	Thr ACC	Ile ATC	Thr ACC	Val GTG	170 Leu CTG
65	Phe TTC	Asn AAC	Thr ACC	Asp GAC	Glu GAG	Glu GAA	Lys AAG	Gly GGT	Ser TCC

**0 121 352**

	180 Phe TTC	Gly GGC	Ser TCG	Arg CGC	Asp GAC	Leu CTG	Ile ATC	Gln CAG	Glu GAA
5	Glu GAA	190 Ala GCC	Lys AAG	Leu CTG	Ala GCC	Asp GAC	Tyr TAC	Val GTG	Leu CTC
10	Ser TCC	Phe TTC	200 Glu GAG	Pro CCC	Thr ACC	Ser AGC	Ala GCA	Gly GGC	Asp GAC
15	Glu GAA	Lys AAA	Leu CTC	210 Ser TCG	Leu CTG	Gly GGC	Thr ACC	Ser TCG	Gly GGC
20	Ile ATC	Ala GCC	Tyr TAC	Val GTG	220 Gln CAG	Val GTC	Asn AAC	Ile ATC	Thr ACC
25	Gly GGC	Lys AAG	Ala GCC	Ser TCG	His CAT	230 Ala GCC	Gly GGC	Ala GCC	Ala GCG
30	Pro CCC	Glu GAG	Leu CTG	Gly GGC	Val GTG	Asn AAC	240 Ala GCG	Leu CTG	Val GTC
35	Glu GAG	Ala GCT	Ser TCC	Asp GAC	Leu CTC	Val GTG	Leu CTG	250 Arg CGC	Thr ACG
40	Met ATG	Asn AAC	Ile ATC	Asp GAC	Asp GAC	Lys AAG	Ala GCG	Lys AAG	260 Asn AAC
45	Leu CTG	Arg CGC	Phe TTC	Asn AAC	Trp TGG	Thr ACC	Ile ATC	Ala GCC	Lys AAG
50	270 Ala GCC	Gly GGC	Asn AAC	Val GTC	Ser TCG	Asn AAC	Ile ATC	Ile ATC	Pro CCC
55	Ala GCC	280 Ser AGC	Ala GCC	Thr ACG	Leu CTG	Asn AAC	Ala GCC	Asp GAC	Val GTG
60	Arg CGC	Tyr TAC	290 Ala GCG	Arg CGC	Asn AAC	Glu GAG	Asp GAC	Phe TTC	Asp GAC
65	Ala GCC	Ala GCC	Met ATG	300 Lys AAG	Thr ACG	Leu CTG	Glu GAA	Glu GAG	Arg CGC
	Ala GCG	Gln CAG	Gln CAG	Lys AAG	310 Lys AAG	Leu CTG	Pro CCC	Glu GAG	Ala GCC
	Asp GAC	Val GTG	Lys AAG	Val GTG	Ile ATC	320 Val GTC	Thr ACG	Arg CGC	Gly GGC

# 0 121 352

	Arg CGC	Pro CCG	Ala GCC	Phe TTC	Asn AAT	Ala GCC	330 Gly GGC	Glu GAA	Gly GGC
5	Gly GGC	Lys AAG	Lys AAG	Leu CTG	Val GTC	Asp GAC	Lys AAG	340 Ala GCG	Val GTG
10	Ala GCC	Tyr TAC	Tyr TAC	Lys AAG	Glu GAA	Ala GCC	Gly GGC	Gly GGC	350 Thr ACG
	Leu CTG	Gly GGC	Val GTG	Glu GAA	Glu GAG	Arg CGC	Thr ACC	Gly GGC	Gly GGC
15	360 Gly GGC	Thr ACC	Asp GAC	Ala GCG	Ala GCC	Tyr TAC	Ala GCC	Ala GCG	Leu CTC
20	Ser TCA	370 Gly GGC	Lys AAG	Pro CCA	Val GTG	Ile ATC	Glu GAG	Ser AGC	Leu CTG
25	Gly GGC	Leu CTG	380 Pro CCG	Gly GGC	Phe TTC	Gly GGC	Tyr TAC	His CAC	Ser AGC
30	Asp GAC	Lys AAG	Ala GCC	390 Glu GAG	Tyr TAC	Val GTG	Asp GAC	Ile ATC	Ser AGC
35	Ala GCG	Ile ATT	Pro CCG	Arg CGC	400 Arg CGC	Leu CTG	Tyr TAC	Met ATG	Ala GCT
	Ala CGC	Arg CGC	Leu CTG	Ile ATC	Met ATG	410 Asp GAT	Leu CTG	Gly GGC	Ala GCC
40	Gly GGC	Lys AAG	TGA	—3'					

8. A recombinant DNA transfer vector according to any preceding claim characterised in that the transfer vector is a plasmid.

45 9. A microorganism transformed by a transfer vector characterised in that the transfer vector is a recombinant DNA transfer vector according to any one of the preceding claims 1 to 8.

10. A microorganism according to claim 9 which is a bacterium of the species *E. coli* or *Pseudomonas* or a yeast of the species *Saccharomyces cerevisiae*.

11. A process for the preparation of a gene product characterised by  
50 (a) culturing a microorganism according to either claim 9 or claim 10 in a culture medium to produce the gene product in the culture medium or the periplasmic space of the microorganism, and  
(b) isolating the gene product from the culture medium or the periplasmic space of the microorganism.

12. A process according to claim 11 characterised in that the gene product is *Pseudomonas* carboxypeptidase G<sub>2</sub>.

13. A process according to claim 11 characterised in that the gene product is a protein or polypeptide other than *Pseudomonas* carboxypeptidase G<sub>2</sub>.

14. A process according to claim 13 characterised in that the gene product is a prokaryotic protein other than *Pseudomonas* carboxypeptidase G<sub>2</sub>.

15. A process according to claim 14 characterised in that the gene product is *E. coli* β-galactosidase.

60

## Patentansprüche

1. Rekombinanter DNA-Transfervektor, enthaltend ein Leitsequenz-polynukleotid stromabwärts von  
65 und in Lesephase mit einem Bakterien- oder Hefepromotor und einer prokaryotischen Ribosomen-

# 0 121 352

Bindestelle und stromaufwärts von und in Lese-Phase mit einem Strukturgen, dadurch gekennzeichnet, daß das Leitsequenz-polynukleotid für ein Signalpolypeptid der Formel I codiert

5 Met-Arg-Pro-Ser-Ile-His-Arg-Thr-  
Ala-Ile-Ala-Ala-Val-Leu-Ala-Thr-  
Ala-Phe-Val-Ala-Gly-Thr.

2. Rekombinanter DNA-Transfervektor nach Anspruch 1, dadurch gekennzeichnet, daß das Leitsequenz-polynukleotid die Formel II aufweist

10 5'— ATG CGC CCA TCC ATC CAC CGC ACA  
GCC ATC GCC GCC GTG CTG GCC ACC II  
GCC TTC GTG GCG GGC ACC —3'.

15 3. Rekombinanter DNA-Transfervektor nach Anspruch 1 oder 2, dadurch gekennzeichnet, daß das Strukturgen für Pseudomonas-Carboxypeptidase G<sub>2</sub> (CPG<sub>2</sub>) codiert.

4. Rekombinanter DNA-Transfervektor nach Anspruch 1 oder 2, dadurch gekennzeichnet, daß das Strukturgen für ein anderes Protein oder Polypeptid codiert als Pseudomonas-Carboxypeptidase G<sub>2</sub>.

5. Rekombinanter DNA-Transfervektor nach Anspruch 4, dadurch gekennzeichnet, daß das Strukturgen für ein anderes prokaryotisches Protein als Pseudomonas-Carboxypeptidase G<sub>2</sub> codiert.

6. Rekombinanter DNA-Transfervektor nach Anspruch 5, dadurch gekennzeichnet, daß das Strukturgen für β-Galactosidase aus E. coli codiert.

7. Rekombinanter DNA-Transfervektor nach Anspruch 3, enthaltend ein Polynukleotid der Formel

25 1 Met Arg Pro Ser Ile His Arg Thr  
5'— ATG CGC CCA TCC ATC CAC CGC ACA  
10 Ala Ile Ala Ala Val Leu Ala Thr Ala  
30 GCC ATC GCC GCC GTG CTG GCC ACC GCC  
20 Phe Val Ala Gly Thr Ala Leu Ala Gln  
35 TTC GTG GCG GGC ACC GCC CTG GCC CAG  
30 Lys Arg Asp Asn Val Leu Phe Gln Ala  
40 AAG CGC GAC AAC GTG CTG TTC CAG GCA  
40 Ala Thr Asp Glu Gln Pro Ala Val Ile  
GCT ACC GAC GAG CAG CCG GCC GTG ATC  
45 Lys Thr Leu Glu Lys 50 Leu Val Asn Ile  
AAG ACG CTG GAG AAG CTG GTC AAC ATC  
50 Glu Thr Gly Thr Gly 60 Asp Ala Glu Gly  
GAG ACC GGC ACC GGT GAC GCC GAG GGC  
55 Ile Ala Ala Ala Gly Asn Phe 70 Leu Glu  
ATC GCC GCT GCG GGC AAC TTC CTC GAG  
Ala Glu Leu Lys Asn Leu Gly Phe 80 Thr  
GCC GAG CTC AAG AAC CTC GGC TTC ACG  
60 Val Thr Arg Ser Lys Ser Ala Gly Gly Leu  
GTC ACG CGA AGC AAG TCG GCC GGC CTG

65



0 121 352

	90	Val	Val	Gly	Asp	Asn	Ile	Val	Gly	Lys
		GTG	GTG	GGC	GAC	AAC	ATC	GTG	GGC	AAG
5		Ile	100	Gly	Arg	Gly	Gly	Lys	Asn	Leu
		ATC	Lys	GGC	CGC	GGC	GGC	AAG	AAC	CTG
				110	Ser	His	Met	Asp	Thr	Val
10		Leu	Leu	Met	TCG	CAC	ATG	GAC	ACC	GTC
		CTG	CTG	ATG						
		Tyr	Leu	Lys	120	Ile	Leu	Ala	Lys	Ala
15		TAC	CTC	AAG	Gly	ATT	CTC	GCG	AAG	GCC
		Pro	Phe	Arg	130		Gly	Asp	Lys	Ala
		CCG	TTC	CGC	Glu		GGC	GAC	AAG	GCC
20				Val	GAA					
		Tyr	Gly	Pro	Gly	Ile	140	Asp	Asp	Lys
		TAC	GGC	CCG	GGC	ATC	Ala	GAC	GAC	AAG
							GCC			
25		Gly	Gly	Asn	Ala	Val	150	His	Thr	
		GGC	GGC	AAC	GCG	GTC	Leu	CAC	ACG	
							CTG			
30		Leu	Lys	Leu	Leu	Lys	Glu	Tyr	160	Val
		CTC	AAG	CTG	CTG	AAG	GAA	TAC	Gly	GTG
									GGC	
35		Arg	Asp	Tyr	Gly	Thr	Ile	Thr	Val	170
		CGC	GAC	TAC	GGC	ACC	ATC	ACC	GTG	Leu
										CTG
		Phe	Asn	Thr	Asp	Glu	Glu	Lys	Gly	Ser
		TTC	AAC	ACC	GAC	GAG	GAA	AAG	GGT	TCC
40										
		180	Gly	Ser	Arg	Asp	Leu	Ile	Gln	Glu
		Phe	GGC	TCG	CGC	GAC	CTG	ATC	CAG	GAA
		TTC								
45		Glu	190	Lys	Leu	Ala	Asp	Tyr	Val	Leu
		GAA	Ala	AAG	CTG	GCC	GAC	TAC	GTG	CTC
			GCC							
50		Ser	Phe	200	Pro	Thr	Ser	Ala	Gly	Asp
		TCC	TTC	Glu	CCC	ACC	AGC	GCA	GGC	GAC
				GAG						
		Glu	Lys	Leu	210	Leu	Gly	Thr	Ser	Gly
55		GAA	AAA	CTC	Ser	CTG	GGC	ACC	TCG	GGC
					TCG					
		Ile	Ala	Tyr	Val	220	Val	Asn	Ile	Thr
		ATC	GCC	TAC	GTG	Gln	GTC	AAC	ATC	ACC
						CAG				
60		Gly	Lys	Ala	Ser	His	230	Gly	Ala	Ala
		GGC	AAG	GCC	TCG	CAT	Ala	GGC	GCC	GCG
							GCC			
65										

0 121 352

	Pro CCC	Glu GAG	Leu CTG	Gly GGC	Val GTG	Asn AAC	240 Ala GCG	Leu CTG	Val GTC
5	Glu GAG	Ala GCT	Ser TCC	Asp GAC	Leu CTC	Val GTG	Leu CTG	250 Arg CGC	Thr ACG
10	Met ATG	Asn AAC	Ile ATC	Asp GAC	Asp GAC	Lys AAG	Ala GCG	Lys AAG	260 Asn AAC
	Leu CTG	Arg CGC	Phe TTC	Asn AAC	Trp TGG	Thr ACC	Ile ATC	Ala GCC	Lys AAG
15	270 Ala GCC	Gly GGC	Asn AAC	Val GTC	Ser TCG	Asn AAC	Ile ATC	Ile ATC	Pro CCC
20	Ala GCC	280 Ser AGC	Ala GCC	Thr ACG	Leu CTG	Asn AAC	Ala GCC	Asp GAC	Val GTG
25	Arg CGC	Tyr TAC	290 Ala GCG	Arg CGC	Asn AAC	Glu GAG	Asp GAC	Phe TTC	Asp GAC
30	Ala GCC	Ala GCC	Met ATG	300 Lys AAG	Thr ACG	Leu CTG	Glu GAA	Glu GAG	Arg CGC
35	Ala GCG	Gln CAG	Gln CAG	Lys AAG	310 Lys AAG	Leu CTG	Pro CCC	Glu GAG	Ala GCC
	Asp GAC	Val GTG	Lys AAG	Val GTG	Ile ATC	320 Val GTC	Thr ACG	Arg CGC	Gly GGC
40	Arg CGC	Pro CCG	Ala GCC	Phe TTC	Asn AAT	Ala GCC	330 Gly GGC	Glu GAA	Gly GGC
45	Gly GGC	Lys AAG	Lys AAG	Leu CTG	Val GTC	Asp GAC	Lys AAG	340 Ala GCG	Val GTG
50	Ala GCC	Tyr TAC	Tyr TAC	Lys AAG	Glu GAA	Ala GCC	Gly GGC	Gly GGC	350 Thr ACG
	Leu CTG	Gly GGC	Val GTG	Glu GAA	Glu GAG	Arg CGC	Thr ACC	Gly GGC	Gly GGC
55	360 Gly GGC	Thr ACC	Asp GAC	Ala GCG	Ala GCC	Tyr TAC	Ala GCC	Ala GCG	Leu CTC
60	Ser TCA	370 Gly GGC	Lys AAG	Pro CCA	Val GTG	Ile ATC	Glu GAG	Ser AGC	Leu CTG
65	Gly GGC	Leu CTG	380 Pro CCG	Gly GGC	Phe TTC	Gly GGC	Tyr TAC	His CAC	Ser AGC

# 0 121 352

	Asp GAC	Lys AAG	Ala GCC	390 Glu GAG	Tyr TAC	Val GTG	Asp GAC	Ile ATC	Ser AGC
5	Ala GCG	Ile ATT	Pro CCG	Arg CGC	400 Arg CGC	Leu CTG	Tyr TAC	Met ATG	Ala GCT
10	Ala CGC	Arg CGC	Leu CTG	Ile ATC	Met ATG	410 Asp GAT	Leu CTG	Gly GGC	Ala GCC
	Gly GGC	Lys AAG	TGA	—3'					

- 15 8. Rekombinanter DNA-Transfervektor nach einem der vorstehenden Ansprüche, dadurch gekennzeichnet, daß der Transfervektor ein Plasmid ist.
9. Durch einen Transfervektor transformierter Mikroorganismus, dadurch gekennzeichnet, daß der Transfervektor ein rekombinanter DNA-Transfervektor nach einem der Ansprüche 1 bis 8 ist.
- 20 10. Mikroorganismus nach Anspruch 9, der ein Bakterium der Gattung *E. coli* oder *Pseudomonas* oder eine Hefe der Gattung *Saccharomyces cerevisiae* ist.
11. Verfahren zur Herstellung eines Genprodukts, gekennzeichnet durch
- (a) Züchten eines Mikroorganismus nach Anspruch 9 oder 10 in einem Kulturmedium zur Herstellung des Genprodukts in Kulturmedium oder im periplasmatischen Raum des Mikroorganismus, und
- 25 (b) Isolieren des Genprodukts aus dem Kulturmedium oder dem periplasmatischen Raum des Mikroorganismus.
12. Verfahren nach Anspruch 11, gekennzeichnet durch Herstellung von *Pseudomonas*-Carboxypeptidase  $G_2$  als Genprodukt.
13. Verfahren nach Anspruch 11, gekennzeichnet durch Herstellung eines anderen Proteins oder
- 30 Polypeptids als *Pseudomonas*-Carboxypeptidase  $G_2$  als Genprodukt.
14. Verfahren nach Anspruch 13, gekennzeichnet durch eines anderen prokaryotischen Proteins als *Pseudomonas*-Carboxypeptidase  $G_2$  als Genprodukt.
15. Verfahren nach Anspruch 14, gekennzeichnet durch  $\beta$ -Galactosidase aus *E. coli* als Genprodukt.

## 35 Revendications

1. Un vecteur de transfert d'ADN recombinant comprenant une séquence guide de polynucléotide en aval et en phase de lecture relativement à un promoteur de bactérie ou de levure et à un site de liaison ribosomique procaryotique et en amont et en phase de lecture relativement à un gène de structure,
- 40 caractérisé en ce que la séquence guide de polynucléotide code pour un polypeptide signal de formule I

Met-Arg-Pro-Ser-Ile-His-Arg-Thr-Ala-Ile-Ala  
Ala-Val-Leu-Ala-Thr-Ala-Phe-Val-Ala-Gly-Thr.

I

- 45 2. Un vecteur de transfert d'ADN recombinant selon la revendication 1, caractérisé en ce que la séquence guide de polynucléotide répond à la formule II

5'	—	ATG	CGC	CCA	TCC	ATC	CAC	CGC	ACA	
		GCC	ATC	GCC	GCC	GTG	CTG	GCC	ACC	II
50		GCC	TTC	GTG	GCG	GGC	ACC	—3'		

3. Un vecteur de transfert d'ADN recombinant selon l'une ou l'autre des revendications 1 ou 2, caractérisé en ce que le gène de structure code pour la carboxypeptidase  $G_2$  de *Pseudomonas* (CPG $_2$ ).
4. Un vecteur de transfert d'ADN recombinant selon l'une ou l'autre des revendications 1 ou 2,
- 55 caractérisé en ce que le gène de structure code pour une protéine ou un polypeptide autre que la carboxypeptidase  $G_2$  de *Pseudomonas*.
5. Un vecteur de transfert d'ADN recombinant selon la revendication 4 caractérisé en ce que le gène de structure code pour une protéine procaryotique autre que la carboxypeptidase  $G_2$  de *Pseudomonas*.
6. Un vecteur de transfert d'ADN recombinant selon la revendication 5 caractérisé en ce que le gène de
- 60 structure code pour la  $\beta$ -galactosidase de *E. coli*.
7. Un vecteur de transfert d'ADN recombinant selon la revendication 3 comprenant un polynucléotide de formule

65

0 121 352

	5'—	1 Met ATG	Arg CGC	Pro CCA	Ser TCC	Ile ATC	His CAC	Arg CGC	Thr ACA
5	Ala GCC	10 Ile ATC	Ala GCC	Ala GCC	Val GTG	Leu CTG	Ala GCC	Thr ACC	Ala GCC
10	Phe TTC	Val GTG	20 Ala GCG	Gly GGC	Thr ACC	Ala GCC	Leu CTG	Ala GCC	Gln CAG
15	Lys AAG	Arg CGC	Asp GAC	30 Asn AAC	Val GTG	Leu CTG	Phe TTC	Gln CAG	Ala GCA
20	Ala GCT	Thr ACC	Asp GAC	Glu GAG	40 Gln CAG	Pro CCG	Ala GCC	Val GTG	Ile ATC
25	Lys AAG	Thr ACG	Leu CTG	Glu GAG	Lys AAG	50 Leu CTG	Val GTC	Asn AAC	Ile ATC
30	Ile ATC	Ala GCC	Ala GCT	Ala GCG	Gly GGC	Asn AAC	Phe TTC	60 Ala GCC	Glu GAG
35	Ala GCC	Glu GAG	Leu CTC	Lys AAG	Asn AAC	Leu CTC	Gly GGC	70 Leu CTC	80 Thr ACG
40	Val GTC	Thr ACG	Arg CGA	Ser AGC	Lys AAG	Ser TCG	Ala GCC	Gly GGC	Leu CTG
45	90 Val GTG	Val GTG	Gly GGC	Asp GAC	Asn AAC	Ile ATC	Val GTG	Gly GGC	Lys AAG
50	Ile ATC	100 Lys AAG	Gly GGC	Arg CGC	Gly GGC	Gly GGC	Lys AAG	Asn AAC	Leu CTG
55	Leu CTG	Leu CTG	110 Met ATG	Ser TCG	His CAC	Met ATG	Asp GAC	Thr ACC	Val GTC
60	Tyr TAC	Leu CTC	Lys AAG	120 Gly GGC	Ile ATT	Leu CTC	Ala GCG	Lys AAG	Ala GCC
65	Pro CCG	Phe TTC	Arg CGC	Val GTC	130 Glu GAA	Gly GGC	Asp GAC	Lys AAG	Ala GCC
	Tyr TAC	Gly GGC	Pro CCG	Gly GGC	Ile ATC	140 Ala GCC	Asp GAC	Asp GAC	Lys AAG

# 0 121 352

	Gly GGC	Gly GGC	Asn AAC	Ala GCG	Val GTC	Ile ATC	150 Leu CTG	His CAC	Thr ACG
5	Leu CTC	Lys AAG	Leu CTG	Leu CTG	Lys AAG	Glu GAA	Tyr TAC	160 Gly GGC	Val GTG
10	Arg CGC	Asp GAC	Tyr TAC	Gly GGC	Thr ACC	Ile ATC	Thr ACC	Val GTG	170 Leu CTG
15	Phe TTC	Asn AAC	Thr ACC	Asp GAC	Glu GAG	Glu GAA	Lys AAG	Gly GGT	Ser TCC
180	Phe TTC	Gly GGC	Ser TCG	Arg CGC	Asp GAC	Leu CTG	Ile ATC	Gln CAG	Glu GAA
20	Glu GAA	190 Ala GCC	Lys AAG	Leu CTG	Ala GCC	Asp GAC	Tyr TAC	Val GTG	Leu CTC
25	Ser TCC	Phe TTC	200 Glu GAG	Pro CCC	Thr ACC	Ser AGC	Ala GCA	Gly GGC	Asp GAC
30	Glu GAA	Lys AAA	Leu CTC	210 Ser TCG	Leu CTG	Gly GGC	Thr ACC	Ser TCG	Gly GGC
35	Ile ATC	Ala GCC	Tyr TAC	Val GTG	220 Gln CAG	Val GTC	Asn AAC	Ile ATC	Thr ACC
	Gly GGC	Lys AAG	Ala GCC	Ser TCG	His CAT	230 Ala GCC	Gly GGC	Ala GCC	Ala GCC
40	Pro CCC	Glu GAG	Leu CTG	Gly GGC	Val GTG	Asn AAC	240 Ala GCG	Leu CTG	Val GTC
45	Glu GAG	Ala GCT	Ser TCC	Asp GAC	Leu CTC	Val GTG	Leu CTG	250 Arg CGC	Thr ACG
50	Met ATG	Asn AAC	Ile ATC	Asp GAC	Asp GAC	Lys AAG	Ala GCG	Lys AAG	260 Asn AAC
	Leu CTG	Arg CGC	Phe TTC	Asn AAC	Trp TGG	Thr ACC	Ile ATC	Ala GCC	Lys AAG
55	270 Ala GCC	Gly GGC	Asn AAC	Val GTC	Ser TCG	Asn AAC	Ile ATC	Ile ATC	Pro CCC
60	Ala GCC	280 Ser AGC	Ala GCC	Thr ACG	Leu CTG	Asn AAC	Ala GCC	Asp GAC	Val GTG
65	Arg CGC	Tyr TAC	290 Ala GCG	Arg CGC	Asn AAC	Glu GAG	Asp GAC	Phe TTC	Asp GAC

# 0 121 352

	Ala GCC	Ala GCC	Met ATG	300 Lys AAG	Thr ACG	Leu CTG	Glu GAA	Glu GAG	Arg GGC
5	Ala GCG	Gln CAG	Gln CAG	Lys AAG	310 Lys AAG	Leu CTG	Pro CCC	Glu GAG	Ala GCC
10	Asp GAC	Val GTG	Lys AAG	Val GTG	Ile ATC	320 Val GTC	Thr ACG	Arg CGC	Gly GGC
15	Arg CGC	Pro CCG	Ala GCC	Phe TTC	Asn AAT	Ala GCC	330 Gly GGC	Glu GAA	Gly GGC
20	Gly GGC	Lys AAG	Lys AAG	Leu CTG	Val GTC	Asp GAC	Lys AAG	340 Ala GCG	Val GTG
	Ala GCC	Tyr TAC	Tyr TAC	Lys AAG	Glu GAA	Ala GCC	Gly GGC	Gly GGC	350 Thr ACG
25	Leu CTG	Gly GGC	Val GTG	Glu GAA	Glu GAG	Arg CGC	Thr ACC	Gly GGC	Gly GGC
30	360 Gly GGC	Thr ACC	Asp GAC	Ala GCG	Ala GCC	Tyr TAC	Ala GCC	Ala GCG	Leu CTC
35	Ser TCA	370 Gly GGC	Lys AAG	Pro CCA	Val GTG	Ile ATC	Glu GAG	Ser AGC	Leu CTG
	Gly GGC	Leu CTG	380 Pro CCG	Gly GGC	Phe TTC	Gly GGC	Tyr TAC	His CAC	Ser AGC
40	Asp GAC	Lys AAG	Ala GCC	390 Glu GAG	Tyr TAC	Val GTG	Asp GAC	Ile ATC	Ser AGC
45	Ala GCG	Ile ATT	Pro CCG	Arg CGC	400 Arg CGC	Leu CTG	Tyr TAC	Met ATG	Ala GCT
50	Ala CGC	Arg CGC	Leu CTG	Ile ATC	Met ATG	410 Asp GAT	Leu CTG	Gly GGC	Ala GCC
	Gly GGC	Lys AAG	TGA	—3'					

55 8. Un vecteur de transfert d'ADN recombinant selon l'une quelconque des revendications précédentes caractérisé en ce que le vecteur de transfert est un plasmide.

9. Un micro-organisme transformé par un vecteur de transfert caractérisé en ce que le vecteur de transfert est un vecteur de transfert d'ADN recombinant selon l'une quelconque des revendications précédentes 1 à 8.

60 10. Un micro-organisme selon la revendication 9 qui est une bactérie de l'espèce *E. coli* ou *Pseudomonas* ou une levure de l'espèce *Saccharomyces cerevisiae*.

11. Un procédé pour la préparation d'un produit génique caractérisé par

(a) la culture d'un micro-organisme selon soit la revendication 9 soit la revendication 10 dans un milieu de culture pour produire le produit génique dans le milieu de culture ou dans l'espace périplasmique du micro-organisme et

(b) l'isolement du produit génique du milieu de culture ou de l'espace périplasmique du micro-organisme.

12. Un procédé selon la revendication 11 caractérisé en ce que le produit génique est la carboxypeptidase G<sub>2</sub> de *Pseudomonas*.

5 13. Un procédé selon la revendication 11 caractérisé en ce que le produit génique est une protéine ou un polypeptide autre que la carboxypeptidase G<sub>2</sub> de *Pseudomonas*.

14. Un procédé selon la revendication 13 caractérisé en ce que le produit génique est une protéine procaryotique autre que la carboxypeptidase G<sub>2</sub> de *Pseudomonas*.

15. Un procédé selon la revendication 14 caractérisé en ce que le produit génique est la  $\beta$ -galactosidase  
10 de *E. coli*.

15

20

25

30

35

40

45

50

55

60

65





Fig1.

RESTRICTION ENZYME MAP OF pNM1


— REPRESENTS pBR322






RESTRICTION ENZYME MAP OF pNM111

*Fig.2.*




RESTRICTION ENZYME MAP OF pNM14

*Fig.3.*



RESTRICTION ENZYME MAP OF pNM21

Fig.4.



RESTRICTION ENZYME MAP OF pNM22

*Fig.5.*





Fig.6.



- pBR 322
- Yeast 2 $\mu$  plasmid
- Yeast chromosomal leu 2 gene
- Pseudomonas carboxypeptidase G2 gene

*Fig.7*

RESTRICTION ENZYME MAP OF pLEC 3